

REAL ALGEBRAIC THREEFOLDS II. MINIMAL MODEL PROGRAM

JÁNOS KOLLÁR

CONTENTS

1. Introduction	1
2. Applications and Speculations	8
2.1. Factorization of Birational Morphisms	8
2.2. Application to the Nash Conjecture	9
2.3. Homology Spheres	10
2.4. Beyond the Nash Conjecture	11
3. The Minimal Model Program over $\mathbb{R}$	12
4. The Topology of Real Points and the MMP	17
5. The Topology of Divisorial Contractions	21
6. The Gateway Method	25
7. Small and Divisor-to-Curve Contractions	30
8. Proof of the Main Theorems	36
9. cAx and cD -type Points	40
10. cA -type Points	44
11. cE -type Points	51
12. Hyperbolic 3-manifolds	57
References	60

1. INTRODUCTION

In real algebraic geometry, one of the main directions of investigation is the topological study of the set of real solutions of algebraic equations. The first general result was proved in [Nash52], and later developed by many others (see [Akbulut-King92] for some recent directions). One of these theorems says that every compact differentiable manifold can be realized as the set of real points of an algebraic variety. [Nash52] posed the problem of obtaining similar results using a restricted class of varieties, for instance rational varieties. For real algebraic surfaces this question was settled in [Comessatti14].

The aim of this series of papers is to utilize the theory of minimal models to investigate this question for real algebraic threefolds. This approach is very similar in spirit to the one employed by [Comessatti14]. (See [Silhol89, Kollár97a] for introductions to real algebraic surfaces from the point of view of the minimal model program.)

For algebraic threefolds over $\mathbb{C}$, the minimal model program (MMP for short) provides a very powerful tool. The method of the program is the following. (See [Kollár87, CKM88] or [Kollár-Mori98] for introductions.)

Starting with a smooth projective 3-fold X , we perform a series of “elementary” birational transformations

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_n =: X^*$$

until we reach a variety X^* whose global structure is “simple”. (Neither the intermediate steps X_i nor the final X^* are uniquely determined by X .) In essence the minimal model program allows us to investigate many questions in two steps: first study the effect of the “elementary” transformations and then consider the “simple” global situation.

In practice both of these steps are frequently rather difficult. For instance, we still do not have a complete list of all possible “elementary” steps, despite repeated attempts to obtain it.

A somewhat unpleasant feature of the theory is that the varieties X_i are not smooth, but have so called terminal singularities. This means that $X_i(\mathbb{R})$ is not necessarily a manifold. In developing the theory of minimal models for real algebraic threefolds, we again have to understand the occurring terminal singularities. This was done in the first paper of this series [Kollár97b].

If X is defined over a field K , then there is a variant of the MMP where the intermediate varieties X_i are also defined over K . I refer to this as the MMP over K . This suggests the following two step approach to understand the topology of $X(\mathbb{R})$:

1. Study the topological effect of the “elementary” transformations.
2. Investigate the topology of $X^*(\mathbb{R})$.

The aim of this paper is to complete the first of these two steps.

I am unable to say much about this question in general. There are serious problems coming from algebraic geometry and also from 3-manifold topology. Some of these are discussed in section 4. My aim is therefore more limited: find reasonable conditions which ensure that the steps of the MMP can be described topologically.

The simplest case to study is contractions $f : X \rightarrow Y$ where X is smooth. Over $\mathbb{C}$ the complete list of such contractions is known [Mori82], and it is not hard to obtain a complete list over $\mathbb{R}$. From

this list one can see that in all such examples where $X(\mathbb{R}) \rightarrow Y(\mathbb{R})$ is complicated, $X(\mathbb{R})$ contains a special surface of nonnegative Euler characteristic. This turns out to be a general pattern, though the proof presented here relies on a laborious case analysis. The precise technical theorem is stated in (1.7).

None of the complicated examples occur if $X(\mathbb{R})$ is orientable, and this yields the following:

Theorem 1.1. *Let X be a smooth, projective, real algebraic 3-fold and X^* the result of the MMP over $\mathbb{R}$. Assume that $X(\mathbb{R})$ is orientable.*

Then the topological normalization $\overline{X^(\mathbb{R})}$ of $X^*(\mathbb{R})$ is a PL-manifold, and $X(\mathbb{R})$ can be obtained from $\overline{X^*(\mathbb{R})}$ by repeated application of the following operations:*

0. *throwing away all isolated points of $\overline{X^*(\mathbb{R})}$,*
1. *taking connected sums of connected components,*
2. *taking connected sum with $S^1 \times S^2$,*
3. *taking connected sum with $\mathbb{RP}^3$.*

Remark 1.2. X^* uniquely determines (1.1.0) and also (1.1.1). The latter can be seen by analyzing real analytic morphisms $h : [0, 1] \rightarrow \overline{X^*(\mathbb{R})}$ where the endpoints map to different connected components of $\overline{X^*(\mathbb{R})}$. In practice this may be quite hard, and it could be easier to work through the MMP backwards.

X^* contains some information about the steps (1.1.2–3), but these are by no means unique. Even if X^* is smooth, both of these steps are possible, as shown by the next example.

Example 1.3. It is well known how to create connected sum with $\mathbb{RP}^3$ algebraically. Let X be a smooth 3-fold over $\mathbb{R}$ and $0 \in X(\mathbb{R})$ a real point. Set $Y = B_0 X$. Then $Y(\mathbb{R}) \sim X(\mathbb{R}) \# \mathbb{RP}^3$. (The connected sum of two nonoriented manifolds is, in general, not unique. It is, however, unique if one of the summands has an automorphism with an isolated fixed point which reverses local orientation there.)

Connected sum with $S^1 \times S^2$ is somewhat harder. Let X be a smooth 3-fold over $\mathbb{R}$ and $D \subset X$ a real curve which has a unique real point $\{0\} = D(\mathbb{R})$. Assume furthermore that near 0 the curve is given by equations $(z = x^2 + y^2 = 0)$. Set $Y_1 = B_D X$. Y_1 has a unique singular point P ; set $Y = B_P Y_1$. It is not hard to see that Y is smooth and $Y(\mathbb{R}) \sim X(\mathbb{R}) \# (S^1 \times S^2)$.

Remark 1.4. As (1.1) already shows, we have to move between topological, PL and differentiable manifolds. In dimension 3 every compact topological 3-manifold carries a unique PL-manifold structure

(cf. [Moise77, Sec. 36]) and also a unique differentiable structure (cf. [Hempel76, p.3]). I mostly use the PL-structure since most algebraic constructions are natural in the PL-category. For instance, $\mathbb{R}^1 \rightarrow \mathbb{R}^2$ given by $t \mapsto (t^2, t^3)$ is a PL-embedding but not a differentiable embedding in the natural differentiable structures.

In dimension 3 the PL-structure behaves very much like a differentiable structure. For instance, let M^3 be a PL 3-manifold, N a compact PL-manifold of dimension 1 or 2 and $g : N \hookrightarrow M$ a PL-embedding. Then a suitable open neighborhood of $g(N)$ is PL-homeomorphic to a real vector bundle over N (cf. [Moise77, Secs. 24 and 26]). (Note that a similar result fails for topological 3-manifolds (cf. [Moise77, Sec. 18]), and it also fails for PL 4-manifolds: take any nontrivial knot in S^3 and suspend it in S^4 .)

1.5 (Surfaces in 3-manifolds). Let M be a PL 3-manifold without boundary, N a compact PL 2-manifold without boundary and $g : N \hookrightarrow M$ a PL-embedding. As we noted above, a neighborhood of N is an $\mathbb{R}$ -bundle over N . $\mathbb{R}$ -bundles over N are classified by group homomorphisms $\rho : \pi_1(N) \rightarrow \{\pm 1\}$. If ρ is trivial then N is 2-sided in M , otherwise it is 1-sided. We also allow self homeomorphisms of N , thus we get the following possibilities when N has nonnegative Euler characteristic:

S^2 : Always 2-sided, many such surfaces in every M^3 .

$\mathbb{RP}^2$: M^3 is not orientable in the 2-sided case. Such manifolds are called $\mathbb{P}^2$ -reducible (cf. [Hempel76, p.88]). In the 1-sided case the boundary of a regular neighborhood is S^2 , thus $M \sim M' \# \mathbb{RP}^3$ for some 3-manifold M' . Most 3-manifolds do not contain any $\mathbb{RP}^2$.

Torus: The 2-sided case occurs in any 3-manifold as the boundary of a regular neighborhood of any S^1 along which M is orientable. There is a unique 1-sided case. For these M is not orientable. Most nonorientable 3-manifolds do not contain 1-sided tori, see section 12.

Klein bottle: M is nonorientable in the 2-sided case. The boundary of a regular neighborhood of any S^1 along which M is nonorientable is such. There are two different 1-sided cases, depending on whether M is orientable near N or not. These are again rare, see section 12.

This shows that there are many 3-manifolds which do not contain $\mathbb{RP}^2$, 1-sided tori or Klein bottles. These correspond to 6 different cases on the above list. It turns out that we need to exclude only 3 of these for our main theorem.

Condition 1.6. Let M be a PL 3-manifold without boundary. Consider the following properties:

1. M does not contain a 2-sided $\mathbb{RP}^2$,
2. M does not contain a 1-sided torus,
3. M does not contain a 1-sided Klein bottle with nonorientable neighborhood.

Failure of any of these properties implies that M is not orientable, but there are many nonorientable 3-manifolds which do satisfy all 3 of the above conditions. For instance, this holds if M is hyperbolic (12.1).

Theorem 1.7. *Let X be a smooth, projective, real algebraic 3-fold and X^* the result of the MMP over $\mathbb{R}$. Assume that $X(\mathbb{R})$ satisfies the 3 conditions (1.6.1–3).*

Then the conclusions of (1.1) hold.

Remark 1.8. It would seem that we also need to allow connected sum with $S^1 \tilde{\times} S^2$ (cf. (5.1)), corresponding to attaching a nonorientable 1–handle. This, however, would give a 1–sided torus which we excluded.

All 3 conditions (1.6.1–3) are necessary for the theorem to hold. My feeling is that essentially nothing can be said without (1.6.1) or (1.6.3). (1.6.2) has a twofold role in the proof. First, it ensures that X is not obtained as a blow up of a smooth 3-fold Y along a curve. This in itself would not be a problem, but it may happen that $Y(\mathbb{R})$ contains a 2-sided $\mathbb{RP}^2$ but $X(\mathbb{R})$ does not. It seems to me that this leads to rather complicated topological questions. Still, a suitable reformulation of the theorem may get around this problem. Second, (1.6.2) is also used to exclude a few singularities on the X_i . These cases are of index 1 and they can be described very explicitly. It should be possible to work with them.

The technical heart of the proof is a listing of the possible singularities that occur in the course of the MMP and a fairly detailed description of the steps of the MMP. The final result is relatively easy to state but the proof is a case-by-case examination.

Theorem 1.9. *Let X be a smooth, projective, real algebraic 3-fold and assume that $X(\mathbb{R})$ satisfies the 3 conditions (1.6.1–3).*

Let X_i be any of the intermediate steps of the MMP over $\mathbb{R}$ starting with X and $0 \in X_i(\mathbb{R})$ a real point. Then a neighborhood of $0 \in X_i$ is real analytically equivalent to one of the following standard forms:

1. (cA_0) Smooth point.
2. $(cA_{>0}^+)$ $(x^2 + y^2 + g_{\geq 2}(z, t) = 0)$, where g is not everywhere negative in a punctured neighborhood of 0.

3. $(cE_6) (x^2 + y^3 + (z^2 + t^2)^2 + yg_{\geq 4}(z, t) + g_{\geq 6}(z, t) = 0)$.

Remark 1.10. The symbol $g_{\geq m}$ denotes a power series of multiplicity at least m .

The name of the cases is explained in [Kollár97b].

The above points of type cE_6 form a codimension 7 family in the space of all cE_6 singularities. They all occur, even if $X(\mathbb{R})$ is orientable. Points of type $cA_{>0}^+$ occur for many choices of g . Section 10 gives an algorithm to decide which cases of g do occur, but I was unable to write the condition in closed form. For the applications this does not seem to matter.

Using [Kollár97b, 4.3, 4.4, 4.9], this immediately implies:

Corollary 1.11. *Notation and assumptions as in (1.9). Then $\overline{X_i(\mathbb{R})} \setminus \{\text{isolated points}\}$ is a compact PL 3-manifold without boundary.* $\square$

The next step is to understand the “elementary” steps of the MMP over $\mathbb{R}$. (1.7) turns out to be a consequence of (1.12). (See (9.1) for the definition of weighted blow-ups.)

Theorem 1.12. *Let X be a smooth, projective, real algebraic 3-fold such that $X(\mathbb{R})$ satisfies the conditions (1.6.1–3).*

Let $f_i : X_i \dashrightarrow X_{i+1}$ be any of the intermediate steps of the MMP over $\mathbb{R}$ starting with X . Then the induced map $f_i : X_i(\mathbb{R}) \rightarrow X_{i+1}(\mathbb{R})$ is everywhere defined and the following is a complete list of possibilities for f_i :

1. ($\mathbb{R}$ -trivial) f_i is an isomorphism in a (Zariski) neighborhood of the set of real points.
2. ($\mathbb{R}$ -small) $f_i : X_i(\mathbb{R}) \rightarrow X_{i+1}(\mathbb{R})$ collapses a 1-complex to points and there are small perturbations $\tilde{f}_i$ of f_i such $\tilde{f}_i : \overline{X_i(\mathbb{R})} \rightarrow \overline{X_{i+1}(\mathbb{R})}$ is a PL-homeomorphism.
3. (smooth point blow up) f_i is the inverse of the blow up of a smooth point $P \in X_{i+1}(\mathbb{R})$.
4. (singular point blow up) f_i is the inverse of a (weighted) blow up of a singular point $P \in X_{i+1}(\mathbb{R})$. There are two cases:
 - (a) $(cA_{>0}^+, \text{mult}_0 g \text{ even})$ Up to real analytic equivalence near P , $X_{i+1} \cong (x^2 + y^2 + g_{\geq 2m}(z, t) = 0)$ where $g_{2m}(z, t) \neq 0$, $m \geq 1$ and X_i is the weighted blow up $B_{(m, m, 1, 1)} X_{i+1}$.
 - (b) $(cA_{>0}^+, \text{mult}_0 g \text{ odd})$ Up to real analytic equivalence near P , $X_{i+1} \cong (x^2 + y^2 + g_{\geq 2m+1}(z, t) = 0)$ where $m \geq 1$, $z^{2m+1} \in g$ and $z^i t^j \notin g$ for $2i + j < 4m + 2$. X_i is the weighted blow up $B_{(2m+1, 2m+1, 2, 1)} X_{i+1}$.

Remark 1.13. The more precise results in sections 9–11 give a description of the various cases when f_i is $\mathbb{R}$ -small (though so far I have not excluded some cases).

The $\mathbb{R}$ -trivial steps do not change anything in a neighborhood of the real points, but it is in these steps that the full complexity of the MMP appears. All the difficulties involving higher index terminal singularities and flips are present, but they always appear in conjugate pairs.

For the topological questions these have no effect, but in other applications of (1.12) this should be taken into account.

Remark 1.14. The lists in (1.9) and (1.12) are fairly short, but I do not see a simple conceptual way of stating the results, let alone proving them by general arguments. The appearance of the singularities of type cE_6 in (1.9) was rather unexpected for me.

The formulations also hide the circumstance that there does not seem to be a single method of excluding all other a priori possible cases. The algebraic method of the proof of (1.9) ends with a much longer list (8.4). The topological method excludes many of these right away, but in a few cases several steps of the MMP need to be analyzed.

1.15 (Method of the proof of (1.12)).

The proof relies on rather extensive computations. The first step is a classification of all 3-dimensional terminal singularities over $\mathbb{R}$ and the study of their topological properties. This was carried out in [Kollár97b]. The next step is to gain a good understanding of the resolutions of these singularities. More precisely, we need to understand the “simplest” exceptional divisors in these resolutions. (Simplicity is measured by the discrepancy, cf. (3.3).) Over $\mathbb{C}$ the first step in this direction is [Markushevich96]. A much more detailed study of such exceptional divisors was completed by [Hayakawa97]. Our main emphasis is over $\mathbb{R}$, and it turns out that there is very little overlap between the computations of [Hayakawa97] and those in sections 9–11. Nonetheless, the basic underlying principles are exactly the same.

Acknowledgments . I thank M. Bestvina, S. Gersten, M. Kapovich and G. Mikhalkin for answering my numerous questions about 3-manifold topology and real algebraic geometry. The existence of cE_6 type points in (1.9) was established with the help of V. Alexeev. I have received helpful comments and questions from A. Bertram, M. Fried, L. Katzarkov and B. Mazur.

Partial financial support was provided by the NSF under grant number DMS-9622394.

2. APPLICATIONS AND SPECULATIONS

2.1. Factorization of Birational Morphisms. Let $f : Y \rightarrow X$ be a birational morphism between smooth and projective varieties. It is a very old problem to factor f as a composite of “elementary” birational morphisms. In dimension 2 this is easy to do: f is the composite of blow ups of points. In dimension 3 and over $\mathbb{C}$, the MMP factors f as a composition of divisorial contractions and flips, but these intermediate steps are rather complicated and not too well understood.

If $f : Y \rightarrow X$ is a birational morphism between smooth and projective threefolds over $\mathbb{R}$, then one would like to get a factorization where the intermediate steps are also defined over $\mathbb{R}$. It turns out that if $Y(\mathbb{R})$ is orientable, the answer is very simple. As with minimal models in general, the intermediate steps involve singular varieties though in this case the real singularities are very mild.

Definition 2.1. A real 3-fold X is said to have a cA_1 singularity at $0 \in X(\mathbb{R})$ if in suitable real analytic coordinates X can be given by an equation $(\pm x^2 \pm y^2 \pm z^2 \pm t^m = 0)$ for a suitable choice of signs and $m \geq 1$.

Theorem 2.2. *Let $f : Y \rightarrow X$ be a birational morphism between smooth and projective threefolds over $\mathbb{R}$. Assume that $Y(\mathbb{R})$ satisfies the conditions (1.6). Then f can be factored as*

$$f : Y = X_n \xrightarrow{f_n} X_{n-1} \rightarrow \cdots \rightarrow X_1 \xrightarrow{f_1} X_0 = X,$$

where each X_i has only cA_1 singularities at real points and the following is a complete list of possibilities for the f_i :

1. (smooth point blow up) f_i is the blow up of a smooth point $P \in X_{i-1}(\mathbb{R})$.
2. (singular point blow up) f_i is the blow up of a singular point $P \in X_{i-1}(\mathbb{R})$.
3. (curve blow up) f_i is the blow up of a real curve $C \subset X_{i-1}$. C has only finitely many real points, X_{i-1} is smooth at each of these and in suitable real analytic coordinates C can be written as $(z = x^2 + y^{2m} = 0)$.
4. ($\mathbb{R}$ -trivial) f_i is an isomorphism in a (Zariski) neighborhood of the set of real points.

Remark 2.3. As in (1.12), it is in the $\mathbb{R}$ -trivial steps that the full complexity of the MMP appears. In particular, the $\mathbb{R}$ -trivial steps may be flips where the flipping curve has no real points.

Proof. For purposes of induction we consider the more general case when X is allowed to have cA_1 -type singularities at real points and terminal singularities at complex points. We assume that X is $\mathbb{Q}$ -factorial (that is, a suitable multiple of every Weil divisor is Cartier). Run the real MMP for Y over X to obtain

$$f : Y = X_n \xrightarrow{f_n} X_{n-1} \dashrightarrow \cdots \dashrightarrow X_1 \xrightarrow{f_1} X_0 = X.$$

The proof is by induction on the number of steps it takes the MMP to reach X .

The last step, $f_1 : X_1 \rightarrow X_0 = X$, is a contraction since we work over X . The possibilities for f_1 are described in (1.12). We are done by induction if f_1 is $\mathbb{R}$ -trivial or a smooth point blow up. Assume that f_1 is a singular point blow up. Since X_0 has only cA_1 points, we are in case (1.12.4a) with $m = 1$. f_1 is the ordinary blow up and by explicit computation we see that X_1 still has only cA_1 singularities.

The case when f_1 is $\mathbb{R}$ -small (1.12.2) needs to be studied in greater detail. f_1 can not be a g -extraction (6.6) since cA_1 type points do not have g -extractions other than the one listed above by (8.4). Thus f_1 is the blow up of a curve $C \subset X_0$. Moreover, X_0 is smooth along $C(\mathbb{R})$ and C is locally planar along $C(\mathbb{R})$ by (7.2). $C(\mathbb{R})$ is finite since f_1 is $\mathbb{R}$ -small.

Pick any point $P \in C(\mathbb{R})$ and assume that C is given by real analytic equations ($z = g(x, y) = 0$). By explicit computation, $B_C X_0$ has a unique singular point with equation ($st - g(x, y) = 0$) which is equivalent to ($u^2 - v^2 - g(x, y) = 0$).

X_1 is an intermediate step of an MMP starting with Y , hence its singularities are among those listed (1.9). Thus g has multiplicity 2 and so it can be written as $\pm x^2 \pm y^r$. Since $(g = 0)$ has only the origin as its real solution, $g = \pm(x^2 + y^{2m})$. $\square$

2.2. Application to the Nash Conjecture. The main conclusion of (1.1) and (1.7) is that if we want to understand the topology of $X(\mathbb{R})$ (say when it is orientable), it is sufficient to study the topology of $X^*(\mathbb{R})$ instead. X^* has various useful properties, depending on the conditions imposed on X .

Consider, for instance, the original Nash question: what happens if X is rational. Since the fifties it has been understood that being rational is a very subtle condition and it is very hard to work with. [KoMiMo92] introduced the much more general notion of being *rationally connected*. A X is rationally connected if two general points of $X(\mathbb{C})$ can be connected by an irreducible rational curve. The lines show that $\mathbb{P}^n$ is rationally connected.

The structure theory of [KoMiMo92] implies that a 3-fold X is rationally connected iff X^* falls in one of 3 classes:

1. (Conic fibrations) There is a morphism (over $\mathbb{R}$) $g : X^* \rightarrow S$ onto a surface such that the general fiber is a conic. Correspondingly there is a morphism $X^*(\mathbb{R}) \rightarrow S(\mathbb{R})$ whose general fiber is S^1 or empty. These cases will be studied in a subsequent paper.
2. (Del Pezzo fibrations) There is a morphism (over $\mathbb{R}$) $g : X^* \rightarrow C$ onto a curve such that the general fiber is a Del Pezzo surface. If $X(\mathbb{R})$ is orientable, then this induces a morphism $X^*(\mathbb{R}) \rightarrow C(\mathbb{R})$ whose general fiber is a torus or a union of some copies of S^2 . These cases will be studied later.
3. (Fano varieties) The anticanonical bundle of X^* is ample. There is a complete list of such varieties if X^* is also smooth [Iskovskikh80]. Even if X^* is known rather explicitly, a topological description of $X^*(\mathbb{R})$ may not be easy. It would be interesting to work out at least some of the cases, for instance hypersurfaces of degree 3 or 4 in $\mathbb{P}^4$. (Mikhalkin pointed out that the degree 3 cases can be understood using the classification of degree 4 real surfaces in $\mathbb{RP}^3$ [Kharlamov76].)

In general it is known that there are only finitely many families of singular Fano varieties in dimension 3 [Kawamata92]. Thus we can get only finitely many different topological types for $X^*(\mathbb{R})$ in this case.

2.3. Homology Spheres. It is interesting to consider if we can get further simplifications of the real MMP if we pose further restrictions on $X(\mathbb{R})$. We may assume, for instance, that $X(\mathbb{R})$ is a homology sphere. This was in fact the assumption I considered first. One can ask if under this assumption $X(\mathbb{R}) \rightarrow X^*(\mathbb{R})$ is a homeomorphism.

Unfortunately this is not the case. Consider for instance the singular real threefold X^* given by affine equation

$$x^2 + y^2 + z^2 + (t - a_0)(t - a_m) \prod_{i=1}^{m-1} (t - a_i)^{2r} = 0,$$

where $a_0 < a_1 < \dots < a_m$ are reals. This has $m - 1$ singular points of the form $x^2 + y^2 + z^2 - u^{2r} = 0$, which can be resolved by r successive blow ups. Resolving all singular points we obtain the 3-fold X . One can easily see that $X(\mathbb{R}) \sim S^3$, but $\overline{X^*(\mathbb{R})}$ is the disjoint union of m copies of S^3 .

One may also study the types of singularities that occur if we pose stronger restrictions on $X(\mathbb{R})$. It seems to me that the best one can get is the following:

Conjecture 2.4. *Let X be a smooth projective 3-fold over $\mathbb{R}$. Assume that $X(\mathbb{R})$ satisfies the conditions (1.6) and $X(\mathbb{R})$ can not be written as a connected sum with $S^1 \times S^2$.*

Let X_i be any of the intermediate steps of the MMP over $\mathbb{R}$ starting with X and $0 \in X_i(\mathbb{R})$ a real point. Then a neighborhood of $0 \in X_i$ is real analytically equivalent to one of the following standard forms:

1. (cA_0) *Smooth point.*
2. $(cA_{>0}^+)$ $(x^2 + y^2 + g_{\geq 2}(z, t) = 0)$, *where $\text{mult}_0 g$ is even and g is not everywhere negative in a punctured neighborhood of 0 .*

In fact, most $cA_{>0}^+$ -type singularities should not occur. It is possible that one can write down a complete list. Also, one can be more precise about how the singular points separate $X_i(\mathbb{R})$.

The results in sections 8–11 come close to proving (2.4), but two points remain unresolved. In order to exclude cE_6 type points, one needs to show that the only possible g -extraction is the one described in (11.6). This should be a feasible computation. The main problem is that in (10.10) I could not exclude certain $\mathbb{R}$ -small contractions. I do not see how to deal with this case.

2.4. Beyond the Nash Conjecture. One can refine the 3-dimensional Nash conjecture in two ways.

First, one can study the topology of $X(\mathbb{R})$ for other classes of real algebraic varieties. The simplest cases may be those whose minimal models admit a natural fibration. This should be very helpful in their topological study. One such class is elliptic threefolds, where we have a morphism $X^* \rightarrow S$ whose general fiber is an elliptic curve. A study of the singular fibers occurring in codimension 1 was completed by [Silhol84].

Another, probably more difficult class are Calabi–Yau 3-folds. It would be very interesting to find some connection between the topology of $X(\mathbb{R})$ and mirror symmetry.

The following question is consistent with the examples that I know:

Question 2.5. *Let X be a smooth projective real 3-fold. Assume that $X(\mathbb{R})$ is hyperbolic. Does this imply that X is of general type?*

One can also start with a 3-manifold M and look for a “simple” real projective 3-fold X such that $X(\mathbb{R}) \sim M$. Ideally one would like to find a solution where certain topological structures on M are reflected by the algebraic properties of X .

There are hyperbolic 3-manifolds which embed into $\mathbb{R}^4$. This implies that they can be realized by real algebraic hypersurfaces in $\mathbb{R}^4$. It would be interesting to find such examples.

The methods of this paper require a very detailed study of the steps of the MMP, which is currently feasible only in dimension 3. It would be, however, interesting to develop some examples in higher dimensions.

Example (1.3) describing connected sum with $S^1 \times S^2$ should have interesting higher dimensional versions. There may be other, more complicated examples as well.

The first steps of the 4-dimensional MMP over $\mathbb{C}$ have been recently classified by [Andreatta-Wisniewski96]. It should be possible to obtain the complete list over $\mathbb{R}$ and to study their topology.

3. THE MINIMAL MODEL PROGRAM OVER $\mathbb{R}$

This section is intended to provide a summary of the MMP over $\mathbb{R}$. More generally, I discuss the MMP over an arbitrary field K of characteristic zero, since there is no difference in the general features. Conjecturally the whole program works in all dimensions but at the moment it is only established in dimensions ≤ 3 .

[Kollár87, Kollár90] provide general introductions. The minimal model program for real algebraic surfaces is explained in detail in [Kollár97a]. For more comprehensive treatments (mostly over $\mathbb{C}$) see [CKM88, Kollár et al.92, Kollár-Mori98].

One of the special features of the 3-dimensional MMP is that we have to work with certain singular varieties in the course of the program.

Definition 3.1. Let X be a normal variety defined over a field K . A *(Weil) divisor* over K is a formal linear combination $D := \sum a_i D_i$ ($a_i \in \mathbb{Z}$) of codimension 1 subvarieties, each defined and irreducible over K . A *$\mathbb{Q}$ -divisor* is defined similarly, except we allow $a_i \in \mathbb{Q}$. A divisor D is called *Cartier* if it is locally definable by one equation and *$\mathbb{Q}$ -Cartier* if mD is Cartier for some $m \in \mathbb{N}$. The smallest such $m > 0$ is called the *index* of D .

We say that X is *factorial* (resp. *$\mathbb{Q}$ -factorial*) if every Weil divisor is Cartier (resp. $\mathbb{Q}$ -Cartier).

A divisor D defined over K is Cartier (resp. $\mathbb{Q}$ -Cartier) iff it is Cartier (resp. $\mathbb{Q}$ -Cartier) after some field extension. However, a variety may be $\mathbb{Q}$ -factorial over K and not $\mathbb{Q}$ -factorial over $\bar{K}$. For instance, the cone $x^2 + y^2 + z^2 - t^2$ is factorial over $\mathbb{R}$ but not over $\mathbb{C}$. (For instance, $(x - \sqrt{-1}y = z - t = 0)$ is not $\mathbb{Q}$ -Cartier.)

Definition 3.2. For a normal variety X , let K_X denote its *canonical class*. K_X is a linear equivalence class of Weil divisors. The corresponding reflexive sheaf $\mathcal{O}_X(K_X)$ is isomorphic to the *dualizing sheaf* ω_X of X .

The index of K_X is called the *index* of X .

Definition 3.3. Let X, Y be normal varieties and $f : Y \rightarrow X$ a birational morphism with exceptional set $\text{Ex}(f)$. Let $E_i \subset \text{Ex}(f)$ be the exceptional divisors. If mK_X is Cartier, then $f^*\mathcal{O}_X(mK_X)$ is defined and there is a natural isomorphism

$$f^*\mathcal{O}_X(mK_X)|_{(Y \setminus \text{Ex}(f))} \cong \mathcal{O}_Y(mK_Y)|_{(Y \setminus \text{Ex}(f))}.$$

Hence there are integers b_i such that

$$\mathcal{O}_Y(mK_Y) \cong f^*\mathcal{O}_X(mK_X)(\sum b_i E_i).$$

Formally divide by m and write this as

$$K_Y \equiv f^*(K_X) + \sum a(E_i, X)E_i, \quad \text{where } a(E_i, X) \in \mathbb{Q}.$$

The rational number $a(E_i, X)$ is called the *discrepancy* of E_i with respect to X .

The closure of $f(E_i) \subset X$ is called the *center* of E_i on X . It is denoted by $\text{center}_X E_i$.

If $f' : Y' \rightarrow X$ is another birational morphism and $E'_i := (f' \circ f^{-1})(E_i) \subset Y'$ is a divisor then $a(E'_i, X) = a(E_i, X)$ and $\text{center}_X E_i = \text{center}_X E'_i$. Thus the discrepancy and the center depend only on the divisor up to birational equivalence, but not on the particular variety where the divisor appears.

Definition 3.4. Let X be a normal variety such that K_X is $\mathbb{Q}$ -Cartier. We say that X is *terminal* (or that it has *terminal singularities*) if for every $f : Y \rightarrow X$, the discrepancy of every exceptional divisor is positive.

The following result makes it feasible to decide if X is terminal or not.

Lemma 3.5. *For a normal variety X the following are equivalent:*

1. X is terminal,
2. $a(E, X) > 0$ for every resolution of singularities $f : Y \rightarrow X$ and for every exceptional divisor $E \subset \text{Ex}(f)$.
3. There is a resolution of singularities $f : Y \rightarrow X$ such that $a(E, X) > 0$ for every exceptional divisor $E \subset \text{Ex}(f)$. $\square$

Example 3.6. It is frequently not too hard to compute discrepancies. Assume for instance that X is a hypersurface defined by $(F(x_1, \dots, x_n) = 0)$. A local generator of $\mathcal{O}_X(K_X)$ is given by any of the forms

$$\eta_i := \frac{1}{\partial F / \partial x_i} dx_1 \wedge \dots \wedge dx_{i-1} \wedge dx_{i+1} \wedge \dots \wedge dx_n.$$

Let $f : Y \rightarrow X$ be a resolution of singularities and $P \in Y$ a point with local coordinates $y_1, \dots, y_{n-1}$. f is given by coordinate functions $x_i = f_i(y_1, \dots, y_{n-1})$ and so we can write

$$\begin{aligned} f^* \eta_n &= f^* \left(\frac{1}{\partial F / \partial x_n} \right) \text{Jac} dy_1 \wedge \dots \wedge dy_{n-1}, \quad \text{where} \\ \text{Jac} &= \text{Jac} \left(\frac{f_1, \dots, f_{n-1}}{x_1, \dots, x_{n-1}} \right) \end{aligned}$$

denotes the determinant of the Jacobian matrix. Hence the discrepancies can be computed as the order of vanishing of the Jacobian minus the order of vanishing of $f^*(\partial F / \partial x_n)$.

If X is smooth then we conclude that $a(E, X) \geq 1$ for every exceptional divisor. Thus smooth varieties are terminal.

Next we define various birational maps which have special role in the MMP.

Definition 3.7. Let X be a variety over K and assume that K_X is $\mathbb{Q}$ -Cartier. A proper morphism $g : X \rightarrow Y$ is called an *extremal contraction* if the following conditions hold:

1. $g_* \mathcal{O}_X = \mathcal{O}_Y$,
2. X is $\mathbb{Q}$ -factorial,
3. Let $C \subset X$ be any irreducible curve such that $g(C) = \text{point}$. Then a $\mathbb{Q}$ -divisor D on X is the pull back of a $\mathbb{Q}$ -Cartier $\mathbb{Q}$ -divisor D' on Y iff $(D \cdot C) = 0$. (Necessarily, $D' = g_*(D)$.)

Definition 3.8. Let $g : X \rightarrow Y$ be an extremal contraction.

We say that g is of *fiber type* if $\dim Y < \dim X$.

We say that g is a *divisorial* contraction if the exceptional set $\text{Ex}(g)$ is the support of $\mathbb{Q}$ -Cartier divisor. In this case $\text{Ex}(g)$ is irreducible over K .

We say that g is a *small* contraction if $\dim \text{Ex}(g) \leq \dim X - 2$.

One can see that every extremal contraction is in one of these 3 groups.

Definition 3.9. A proper morphism $f : X \rightarrow Y$ is called *K_X -negative* if $-K_X$ is f -ample.

Definition 3.10. Let $f : X \rightarrow Y$ be a small K_X -negative extremal contraction. A variety X^+ together with a proper birational morphism $f^+ : X^+ \rightarrow Y$ is called a *flip* of f if

1. K_{X^+} is $\mathbb{Q}$ -Cartier,
2. K_{X^+} is f^+ -ample, and
3. the exceptional set $\text{Ex}(f^+)$ has codimension at least two in X^+ .

By a slight abuse of terminology, the rational map $\phi : X \dashrightarrow X^+$ is also called a flip. A flip gives the following diagram:

$$\begin{array}{ccc} X & \xrightarrow{\phi} & X^+ \\ -K_X \text{ is } f\text{-ample} & \searrow \swarrow & K_{X^+} \text{ is } f^+\text{-ample} \\ & Y & \end{array}$$

It is not hard to see that a flip is unique and the main question is its existence.

We are ready to state the 3-dimensional MMP over an arbitrary field:

Theorem 3.11 (MMP over K). *Let X be a smooth projective 3-fold defined over a field K (of characteristic zero). Then there is a sequence*

$$X = X_0 \xrightarrow{f_0} X_1 \dashrightarrow \cdots \dashrightarrow X_i \xrightarrow{f_i} X_{i+1} \dashrightarrow \cdots \xrightarrow{f_{n-1}} X_n =: X^*$$

with the following properties

1. Each X_i is a terminal projective 3-fold over K which is $\mathbb{Q}$ -factorial over K .
2. Each f_i is either a K_X -negative divisorial extremal contraction or the flip of a K_X -negative small extremal contraction.
3. One of the following holds for X^* :
 - (a) K_{X^*} is nef (that is $(C \cdot K_{X^*}) \geq 0$ for any curve $C \subset X^*$), or
 - (b) there is a fiber type extremal contraction $X^* \rightarrow Z$.

Remark 3.12. For the purposes of this paper one can handle the MMP as a black box. It is sufficient to know that it works, but I will use very few of its finer properties. In particular, there is no need to know anything about flips beyond believing their existence.

The rest of the section is devoted to explicitly stating all further results from minimal model theory that I use later. The most significant among these is the classification of terminal 3-fold singularities over nonclosed fields, established in [Kollár97b].

Notation 3.13. For a field K let $K[[x_1, \dots, x_n]]$ denote the ring of formal power series in n variables over K . For $K = \mathbb{R}$ or $K = \mathbb{C}$, let $K\{x_1, \dots, x_n\}$ denote the ring of those formal power series which converge in some neighborhood of the origin.

For a power series F , F_d denotes the degree d homogeneous part. The multiplicity, denoted by $\text{mult}_0 F$, is the smallest d such that $F_d \neq 0$. If we write a power series as $F_{\geq d}$ then it is assumed that its multiplicity is at least d .

For $F \in \mathbb{R}\{x_1, \dots, x_n\}$ let $(F = 0)$ denote the germ of its zero set in $\mathbb{C}^n$ with its natural real structure. I always think of it as a complex analytic germ with a real structure and not just as a real analytic germ in $\mathbb{R}^n$.

$(F = 0)/\frac{1}{n}(a, b, c, d)$ means the following. Define a $\mathbb{Z}_n$ -grading of $\mathbb{C}\{x, y, z, t\}$ by $x \mapsto a, y \mapsto b, z \mapsto c, t \mapsto d$. If F is graded homogeneous, then $(F = 0)/\frac{1}{n}(a, b, c, d)$ denotes the germ whose ring of holomorphic functions is the ring of grade zero elements of $\mathbb{C}\{x, y, z, t\}/(F)$.

If $(F = 0)$ is terminal then n coincides with the index (3.2) of the singularity.

Example 3.14. In case $X = (x^2 + y^2 + z^2 + t^2 = 0)/\frac{1}{2}(1, 1, 1, 0)$ the ring is

$$\mathcal{O}_X = \mathbb{C}\{x^2, y^2, z^2, t, xy, yz, zx\}/(x^2 + y^2 + z^2 + t^2),$$

with the natural real structure.

X can also be realized as the image of the hypersurface $(x^2 + y^2 + z^2 + t^2 = 0)$ under the map

$$\phi : \mathbb{C}^4 \rightarrow \mathbb{C}^7 : (x, y, z, t) \mapsto (x^2, y^2, z^2, t, xy, yz, zx),$$

which has degree 2 over its image.

Although $(x^2 + y^2 + z^2 + t^2 = 0)$ has only the origin as its real solution, X has plenty of real points. Indeed, any real solution of $x^2 + y^2 + z^2 - t^2 = 0$ gives a *real* point $P = \phi(\sqrt{-1}x, \sqrt{-1}y, \sqrt{-1}z, t) \in X(\mathbb{R})$. $\phi^{-1}(P)$ is a pair of conjugate points on the hypersurface $(x^2 + y^2 + z^2 + t^2 = 0)$. All the real elements of $\mathcal{O}_X$ take up real values at P .

This way we see that $X(\mathbb{R})$ is a cone over 2 copies of $\mathbb{RP}^2$.

The following is a summary of the classification of terminal singularities obtained in [Kollár97b]. As it turns out, the classification closely follows the earlier results over algebraically closed fields. The choice of the subdivision into cases is dictated by the needs of the proof in sections 9–11, rather than the internal logic of the classification.

Theorem 3.15. *Let X be a real algebraic or analytic 3-fold and $0 \in X(\mathbb{R})$ a real point. Then X has a terminal singularity at 0 iff a neighborhood of $0 \in X$ is real analytically equivalent to one of the following:*

name	equation
cA_0	$(t = 0)$
cA_1	$(x^2 + y^2 \pm z^2 \pm t^m = 0)$
$cA_{>1}^+$	$(x^2 + y^2 + g_{\geq 3}(z, t) = 0)$
$cA_{>1}^-$	$(x^2 - y^2 + g_{\geq 3}(z, t) = 0)$
cD_4	$(x^2 + f_{\geq 3}(y, z, t) = 0)$, where $f_3 \neq l_1^2 l_2$ for linear forms l_i
$cD_{>4}$	$(x^2 + y^2 z + f_{\geq 4}(y, z, t) = 0)$,
cE_6	$(x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 4}(z, t) = 0)$, where $h_4 \neq 0$
cE_7	$(x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 5}(z, t) = 0)$, where $g_3 \neq 0$
cE_8	$(x^2 + y^3 + yg_{\geq 4}(z, t) + h_{\geq 5}(z, t) = 0)$, where $h_5 \neq 0$
cA_0/n	$(t = 0)/\frac{1}{n}(r, -r, 1, 0)$ where $n \geq 2$ and $(n, r) = 1$
$cA_1/2$	$(x^2 + y^2 \pm z^n \pm t^m = 0)/\frac{1}{2}(1, 1, 1, 0)$ where $\min\{n, m\} = 2$
$cA_{>1}^+/2$	$(x^2 + y^2 + f_{\geq 3}(z, t) = 0)/\frac{1}{2}(1, 1, 1, 0)$
$cA_{>1}^-/2$	$(x^2 - y^2 + f_{\geq 3}(z, t) = 0)/\frac{1}{2}(1, 1, 1, 0)$
cA/n	$(xy + f(z, t) = 0)/\frac{1}{n}(r, -r, 1, 0)$ where $n \geq 3$ and $(n, r) = 1$
$cAx/2$	$(x^2 \pm y^2 + f_{\geq 4}(z, t) = 0)/\frac{1}{2}(0, 1, 1, 1)$
$cAx/4$	$(x^2 \pm y^2 + f_{\geq 2}(z, t) = 0)/\frac{1}{4}(1, 3, 1, 2)$ where $f_2(0, 1) = 0$
$cD/2$	$(x^2 + f_{\geq 3}(y, z, t) = 0)/\frac{1}{2}(1, 1, 0, 1)$
$cD/3$	$(x^2 + f_{\geq 3}(y, z, t) = 0)/\frac{1}{3}(0, 1, 1, 2)$ where $f_3(0, 0, 1) \neq 0$
$cE/2$	$(x^2 + y^3 + f_{\geq 4}(y, z, t) = 0)/\frac{1}{2}(1, 0, 1, 1)$

4. THE TOPOLOGY OF REAL POINTS AND THE MMP

Starting with a projective variety X over $\mathbb{R}$, let us run the MMP over $\mathbb{R}$. We obtain a sequence of birational maps

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_i \xrightarrow{f_i} X_{i+1} \dashrightarrow \cdots \dashrightarrow X^*.$$

These in turn induce (not necessarily everywhere defined) maps between the sets of real points

$$X(\mathbb{R}) = X_0(\mathbb{R}) \dashrightarrow \cdots \dashrightarrow X_i(\mathbb{R}) \xrightarrow{f_i} X_{i+1}(\mathbb{R}) \dashrightarrow \cdots \dashrightarrow X^*(\mathbb{R}).$$

Our aim is to see if there is a way of describing $X(\mathbb{R})$ in terms of $X^*(\mathbb{R})$ and a local description of the maps $X_i(\mathbb{R}) \dashrightarrow X_{i+1}(\mathbb{R})$ in a neighborhood of their exceptional sets.

Proposition 4.1. *Every step f_i of the MMP over $\mathbb{R}$ is among the following five:*

1. (divisor-to-point) f_i contracts a geometrically irreducible divisor $E_i \subset X_i$ to a point $P_{i+1} \in X_{i+1}(\mathbb{R})$.

2. (*divisor-to-curve*) f_i contracts a geometrically irreducible divisor $E_i \subset X_i$ to a real curve $C_{i+1} \subset X_{i+1}$.
3. ($\mathbb{R}$ -small) $f_i : X_i(\mathbb{R}) \rightarrow X_{i+1}(\mathbb{R})$ collapses a 1-complex to points and is a homeomorphism elsewhere.
4. (*flip*) f_i is the flip of a curve $C_i \subset X_i$.
5. ($\mathbb{R}$ -trivial) f_i is an isomorphism in a (Zariski) neighborhood of the set of real points.

Proof. If f_i is a flip then we have case (4). Thus we may assume that f_i is the contraction of a divisor $E_i \subset X_i$ and E_i is irreducible over $\mathbb{R}$. If E_i is irreducible over $\mathbb{C}$ then we have one of the cases (1–2). If E_i is reducible over $\mathbb{C}$ then $E_i(\mathbb{R})$ is a 1-complex by (4.2) and so we are in case (3).

Any of the above cases can also be of type (5). $\square$

Lemma 4.2. *Let X be an n -dimensional scheme over $\mathbb{R}$ (that is, an algebraic variety possibly with several irreducible components and with singularities). Assume that if $X_i \subset X$ is any $\mathbb{R}$ -irreducible component then X_i is reducible over $\mathbb{C}$. Then $X(\mathbb{R}) = (\text{Sing } X)(\mathbb{R})$, that is, every real point is singular. In particular, $\dim X(\mathbb{R}) \leq n - 1$.*

Proof. Assume that $P \in X(\mathbb{R})$ is a smooth real point. Then P lies on a unique irreducible component $Y \subset X_{\mathbb{C}}$, thus Y is invariant under complex conjugation. So Y is an irreducible real component which stays irreducible over $\mathbb{C}$, a contradiction. $\square$

Each of the 5 steps $X_i(\mathbb{R}) \dashrightarrow X_{i+1}(\mathbb{R})$ have different topological behaviour. The following informal discussion intends to emphasize their main features.

4.3 (Divisor-to-point). Let $M = X_i(\mathbb{R})$ be a 3-complex (with only finitely many singular points) and $F = E_i(\mathbb{R}) \subset M$ a 2-complex. We collapse F to a point:

$$\begin{array}{ccc} F & \subset & M \\ \downarrow & & \downarrow f \\ P & \in & N. \end{array}$$

In practice we are frequently able to describe a regular neighborhood $F \subset U \subset M$ (this is a local datum) and by assumption we know a regular neighborhood $P \in V \subset N$. Thus we see that M is obtained from U and $N \setminus \text{Int } V$ by gluing them together along the boundaries ∂U and ∂V .

The gluing is determined by a PL-homeomorphism $\phi : \partial U \rightarrow \partial V$. Thus, besides knowing U and N , we also need to know ϕ up to PL-isotopy. If one of the connected components of ∂U has genus at least 2, this is a very hard problem. In fact, as the example of Heegard

splittings shows (cf. [Hempel76, Ch.2]), the choice of ϕ is usually the most significant information. Unfortunately, ϕ can be described only in terms of global data.

If ∂U is a union of m copies of S^2 , then ϕ is classified by an element of the symmetric group on m elements (which S^2 maps where) and a sign for each S^2 (describing whether the map is orientation preserving or reversing on that S^2). Hence, knowing U and N , we can determine M up to finite ambiguity.

In many cases U is so simple that different choices of ϕ give the same M , giving even fewer possibilities for M .

If $P \in N$ is an isolated singular point, then ∂V is a union of spheres iff $\bar{N}$ (the topological normalization of N) is a manifold.

The situation is similarly simple if ∂U is a union of copies of $\mathbb{RP}^2$ and of S^2 , and still manageable if ∂U also contains tori and Klein bottles. For us these more general cases do not come up.

4.4 (Divisor-to-curve). This time we construct it bottom up. Assume for simplicity that $N = X_{i+1}(\mathbb{R})$ is a 3-manifold and $L = C_{i+1}(\mathbb{R}) \subset N$ a link. The projectivized normal bundle is an S^1 -bundle $S \rightarrow L$. The blow up of L in N replaces L by S to obtain:

$$\begin{array}{ccc} S & \subset & M \\ \downarrow & & \downarrow f \\ L & \in & N. \end{array}$$

(In general N may have finitely many singular points and L is only a 1-complex, but I believe that a similar description is possible in all cases.)

Here M is uniquely determined, once we know N and L . By assumption we know N but L is a free choice. The Jaco–Johannson–Shalen decomposition (cf. [Scott83, p.483]) shows that in most cases $B_L N$ determines $M \setminus L$. Thus the description of all possible $B_L N$ is essentially equivalent to the description of all links.

For us L has to come from an algebraic curve, thus we are led to the question: Which links in a real algebraic 3-fold can be realized by algebraic curves? In some cases every link is realized (cf. [Akbulut-King81]), thus we again run into a hard topological problem.

So M can be described in terms of N , though the answer depends on the choice of a link, which is a very complicated object.

4.5 ($\mathbb{R}$ -small contraction). $N = X_{i+1}(\mathbb{R})$ is obtained from $M = X_i(\mathbb{R})$ by collapsing a 1-complex $C = (\text{Sing } E_i)(\mathbb{R}) = E_i(\mathbb{R})$ to a point:

$$\begin{array}{ccc} C & \subset & M \\ \downarrow & & \downarrow f \\ P & \in & N. \end{array}$$

If the normalizations $\bar{M}$ and $\bar{N}$ are manifolds, then we see in (5.3) that a suitable small perturbation of f is a homeomorphism between $\bar{M}$ and $\bar{N}$. Thus this step (which is actually more complicated from the point of view of algebraic geometry than the previous two cases) is easy to analyze topologically.

4.6 (Flip). Assume for simplicity that $M = X_i(\mathbb{R})$ is an orientable 3-manifold and $C(\mathbb{R}) \sim S^1$. $N = X_{i+1}(\mathbb{R})$ is obtained from M by a surgery along S^1 . The boundary of a regular neighborhood of S^1 is $S^1 \times S^1$, and the surgery is determined by a diffeomorphism of $S^1 \times S^1$ up to isotopy. These are classified by $SL(2, \mathbb{Z})$. (In general M may have finitely many singular points and $C(\mathbb{R})$ is a 1-complex, but I believe that a similar description is possible in all cases.)

A complete classification of flips is known [Kollár-Mori92], thus it should be possible to compute the resulting diffeomorphism of $S^1 \times S^1$.

Here again we run into a global problem. $S^1 \subset M$ may be very knotted, and the result of the surgery depends mostly on the knot $S^1 \subset M$. The usual descriptions of flips characterize a complex analytic neighborhood of C , thus they say nothing about how its real part is knotted. From the point of view of algebraic geometry, this is a global invariant.

We have the additional problem that flipping curves are rigid objects, thus we can not hope to get a flipping curve by approximating a real curve algebraically. Furthermore, it is very hard to determine which curves are obtained by a flip. (Even if Z is a smooth complex 3-fold and $C \subset Z$ a smooth $\mathbb{CP}^1$, I know of no practical way of determining if $C \subset Z$ is obtained as a result of a flip.)

4.7 (Conclusion). Start with a projective 3-fold X over $\mathbb{R}$ and run the MMP over $\mathbb{R}$:

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_i \xrightarrow{f_i} X_{i+1} \dashrightarrow \cdots \dashrightarrow X^*.$$

If we would like to understand the topology of $X(\mathbb{R})$ in terms of $X^*(\mathbb{R})$, then we have to ensure that the MMP has the following properties:

1. $\overline{X_i(\mathbb{R})}$ is a manifold for every i .

2. Each f_i is either $\mathbb{R}$ -trivial or $\mathbb{R}$ -small or a divisor-to-point contraction.

(1.12) asserts that both of these conditions can be satisfied by imposing certain mild conditions on the topology of $X(\mathbb{R})$.

5. THE TOPOLOGY OF DIVISORIAL CONTRACTIONS

The aim of this section is to describe some examples where the change of the topology of a real algebraic variety under a divisorial contraction can be readily understood by topological methods.

Notation 5.1. The disjoint union of two topological spaces is denoted by $M \uplus N$. Direct product is denoted by $M \times N$. The unique nontrivial S^2 -bundle over S^1 is denoted by $S^1 \tilde{\times} S^2$. This is obtained from $[0, 1] \times S^2$ by indentifying the 2 ends via an orientation reversing homeomorphism.

Homeomorphism of two topological spaces is denoted by $M \sim N$.

We start with the study of $\mathbb{R}$ -small contractions:

Lemma 5.2. *Let $f : M \rightarrow N$ be a proper PL-map between PL-manifolds of dimension $n \geq 3$. Assume that there is a 1-complex $C \subset M$ and a finite set of points $P \subset N$ such that $f : M \setminus C \rightarrow N \setminus P$ is a PL-homeomorphism.*

Then M and N are PL-homeomorphic (by a small perturbation of f).

Proof. If C is collapsible to points, then a regular neighborhood of C in M is a union of disjoint n -cells [Hempel76, 1.8] and we are done.

In order to see that C is collapsible to points, we may assume that P is a point and $N = S^n$. Thus M is also a compact PL-manifold. M is orientable outside the codimension ≥ 2 subset C , hence it is orientable. Consider the exact homology sequences

$$\begin{array}{ccccccc} H_i(C) & \rightarrow & H_i(M) & \rightarrow & H_i(M, C) & \rightarrow & H_{i-1}(C) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H_i(P) & \rightarrow & H_i(S^n) & \rightarrow & H_i(S^n, P) & \rightarrow & H_{i-1}(P) \end{array}$$

We compute $H_i(M, C) = H_i(S^n, P)$ from the second sequence and substitute into the first to obtain that

$$H_1(C) \cong H_1(M), \quad \text{and} \quad 0 = H_{n-1}(C) \cong H_{n-1}(M).$$

By Poincaré duality we conclude that $H_1(C) = 0$, thus C is contractible. $\square$

Corollary 5.3. *Let $f : X \rightarrow Y$ be a morphism a n -dimensional real algebraic varieties, $n \geq 3$. Assume that*

1. $\overline{X(\mathbb{R})} = M \uplus R$ and $\overline{Y(\mathbb{R})} = N \uplus R'$ where M, N are PL-manifolds and $\dim R, \dim R' < n$.
2. f induces an isomorphism $R \cong R'$.
3. $\text{Ex}(f)(\mathbb{R})$ is a 1-complex.

Then $\overline{X(\mathbb{R})}$ is PL-homeomorphic to $\overline{Y(\mathbb{R})}$.

Proof. Set $C = \text{Ex}(f)(\mathbb{R})$ and $\bar{C} \subset M \subset \overline{X(\mathbb{R})}$ its preimage. Since $\bar{C}$ has dimension 1, there is a one-to-one correspondence between the connected components of $\bar{C}$ and the connected components of the boundary of a regular neighborhood of $\bar{C}$. Hence f lifts to a morphism $\bar{f} : \overline{X(\mathbb{R})} \rightarrow \overline{Y(\mathbb{R})}$. Thus (5.2) implies (5.3), and the homeomorphism is given by a small perturbation of $\bar{f}$. $\square$

Remark 5.4. A real algebraic curve is a union of copies of S^1 . The proof of (5.2) shows that the preimage of $\text{Ex}(f)(\mathbb{R})$ in $\overline{X(\mathbb{R})}$ is contractible. Thus if f itself is not a homeomorphism, then $X(\mathbb{R})$ is not a manifold.

Next we look at divisor-to-point contractions.

Proposition 5.5. *Let M be a 3-dimensional PL-manifold and $F \subset M$ a connected 2-complex with only finitely many singular points. Let $F \subset \text{Int } U \subset M$ be a regular neighborhood of F . Then*

$$\dim H_1(\bar{F}, \mathbb{Q}) \leq \dim H_1(\partial U, \mathbb{Q}),$$

and strict inequality holds unless every connected component of $\bar{F}$ is one of the following:

1. S^2 or $\mathbb{RP}^2$,
2. a one-sided $S^1 \times S^1$,
3. a one-sided Klein bottle whose neighborhood is not orientable.

Proof. Let F be a compact 2-complex with only finitely many singular points. Its normalization $\bar{F}$ can be written as $F^{(2)} \uplus F^{(1)}$ where $F^{(2)}$ is a compact 2-manifold and a $F^{(1)}$ is a 1-complex.

Pick a point $P \in F$ whose link in F consists of at least 2 circles. Locally F looks like the cone over parallel plane sections $(z = a_i) \cap (x^2 + y^2 + z^2 = 1)$ of the unit sphere in $\mathbb{R}^3$ (plus a few 1-cells). By a homotopy we can replace this by the parallel plane sections of the unit ball $(z = a_i) \cap (x^2 + y^2 + z^2 \leq 1)$ and add the interval $[\min_i \{a_i\}, \max_i \{a_i\}]$ on the z -axis. This does not change the boundary of the regular neighborhood. Thus we may assume that $F^{(2)} \rightarrow M$ is an embedding.

Let us take a point or a 1-cell e in $F^{(1)}$. If e does not intersect the rest of F , then a regular neighborhood of e is a 3-cell. e can be deleted from F without changing the inequality.

If e intersects the rest of F in one endpoint only, then we can delete e from F without changing the regular neighborhood.

If e intersects the rest of F at both endpoints, then removing e creates a new 2-complex F' , and $F^{(2)} = F'^{(2)}$. Let $F' \subset \text{Int } U'$ be its regular neighborhood. ∂U is obtained from $\partial U'$ by attaching a handle $[0, 1] \times S^1$. Thus $H_1(\partial U) \geq H_1(\partial U')$, and it is sufficient to verify our inequality for F' .

At the end we are reduced to the situation when F is the disjoint union of embedded 2-manifolds, and it is sufficient to check the inequality for each connected component of F separately. $\partial U \rightarrow F$ is a 2 sheeted cover, thus $H_1(\partial U) \geq H_1(F)$ with equality only if $F \sim S^2, F \sim \mathbb{RP}^2, F \sim S^1 \times S^1 \sim \partial U$ or F and ∂U are both Klein bottles. $\square$

Proposition 5.6. *Let M be a 3-dimensional PL-manifold and $F \subset M$ a compact 2-complex with only finitely many singular points. Let $0 \in N$ be obtained from M by collapsing F to a point. Assume that $\bar{N}$ is a 3-manifold. Then M can be obtained from $\bar{N}$ by repeated application of the following operations:*

1. taking connected sums of connected components,
2. taking connected sum with $S^1 \times S^2$,
3. taking connected sum with $S^1 \tilde{\times} S^2$, or
4. taking connected sum with $\mathbb{RP}^3$.

Proof. We use the notation of (5.5) and of its proof. Let $F \subset \text{Int } U \subset M$ and $0 \in \text{Int } V \subset N$ be regular neighborhoods such that $U = f^{-1}(V)$. Then $\partial U = \partial V$. Since $\bar{N}$ is a manifold, this implies that ∂U is a union of 2-spheres. We also see that $\bar{N}$ is obtained from $M \setminus \text{Int } U$ by attaching a 3-ball to each S^2 in ∂U .

As in the proof of (5.5) we may assume that $F^{(2)} \rightarrow M$ is an embedding.

If e is a point or a 1-cell in $F^{(1)}$ which intersects the rest of F in zero or one point only, then we can delete e from F .

If e intersects the rest of F at both endpoints, then removing e creates a new 2-complex F' such that $\bar{F} = \bar{F}'$. Let $F' \subset \text{Int } U'$ be its regular neighborhood. ∂U is obtained from $\partial U'$ by attaching a handle $[0, 1] \times S^1$. The two ends $\{0\} \times S^1$ and $\{1\} \times S^1$ can not attach to the same connected component of $\partial U'$ since that would create a torus or a Klein bottle in ∂U . Thus $\partial U'$ has one more copies of S^2 than ∂U .

$\bar{N}$ is obtained from $\bar{N}'$ by collapsing the image of e to a point, hence $\bar{N}$ and $\bar{N}'$ are homeomorphic by (5.2).

At the end we are reduced to the situation when F is the disjoint union of embedded copies of S^2 and $\mathbb{RP}^2$. An S^2 is necessarily 2-sided. Removing it from F corresponds to taking connected sums of connected components (if S^2 separates M) or to taking connected sum with $S^1 \times S^2$ or $S^1 \tilde{\times} S^2$ (if S^2 does not separate M) (cf. [Hempel76, Chap. 3]).

If $\mathbb{RP}^2$ is 2-sided, then the boundary of its regular neighborhood consists of two copies of $\mathbb{RP}^2$, so this can not happen. A 1-sided $\mathbb{RP}^2$ corresponds to taking connected sum with $\mathbb{RP}^3$. $\square$

Corollary 5.7. *Let $f : X \rightarrow Y$ be a morphism a real algebraic 3-folds. Assume that*

1. $\overline{X(\mathbb{R})}$ and $\overline{Y(\mathbb{R})}$ are PL-manifolds, and
2. $\text{Ex}(f)$ is a geometrically irreducible normal surface which is contracted to a point.

Then $\overline{X(\mathbb{R})}$ can be obtained from $\overline{Y(\mathbb{R})}$ by repeated application of the following operations:

3. removing an isolated point from $\overline{Y(\mathbb{R})}$,
4. taking connected sums of connected components,
5. taking connected sum with $S^1 \times S^2$,
6. taking connected sum with $S^1 \tilde{\times} S^2$, or
7. taking connected sum with $\mathbb{RP}^3$.

Proof. If $\text{Ex}(f)(\mathbb{R}) = \emptyset$ then the image of $\text{Ex}(f)$ is an isolated real point of $Y(\mathbb{R})$ which has to be thrown away to obtain $X(\mathbb{R})$. If $\text{Ex}(f)(\mathbb{R}) \neq \emptyset$, then isolated points of $X(\mathbb{R})$ correspond to isolated points of $Y(\mathbb{R})$, hence they can be ignored.

Let M be the topological normalization of $X(\mathbb{R}) \setminus (\text{isolated points})$, N the topological normalization of $Y(\mathbb{R}) \setminus (\text{isolated points})$ and F the preimage of $\text{Ex}(f)(\mathbb{R})$ in M . F is a 2-complex with isolated singularities since $\text{Ex}(f)$ is normal.

Thus (5.7) follows from (5.6). $\square$

Complement 5.8. *It is worthwhile to note that condition (5.7.2) can be weakened to:*

- 2' $\text{Ex}(f)$ contains a unique geometrically irreducible surface S . S has only isolated singularities and S is contracted to a point by f .

It would be very useful to have a version of (5.7) which works if $\text{Ex}(f)$ is an irreducible but nonnormal surface.

In the topological version (5.6) essentially nothing can be said if F is allowed to become an arbitrary compact 2-complex. For instance, let M be an arbitrary compact 3-manifold and F the 2-skeleton of a

triangulation of M . Then $\bar{N}$ is the union of some copies of S^3 (one for each 3-simplex).

This example usually can not arise as the real points of an algebraic surface, but it is not hard to modify this example by approximating each simplex with a sphere to get the following. (This is not used in the sequel and so no proof is given here.)

Proposition 5.9. *Let M be a compact differentiable manifold of dimension n . Then there is a smooth real algebraic variety X and a morphism $f : X \rightarrow Y$ with the following properties:*

1. $X(\mathbb{R}) \sim M$,
2. $\bar{Y}(\mathbb{R})$ is a disjoint union of copies of S^n ,
3. $\text{Ex}(f)$ is a geometrically irreducible divisor and $\text{Ex}(f)(\mathbb{R})$ is a union of copies of S^{n-1} intersecting transversally.

6. THE GATEWAY METHOD

At the beginnings of the MMP, divisorial contractions were considered to be the easily understandable part of the program and flips the hard part. Lately, however, more and more questions require a detailed understanding of all the steps of the MMP. A fairly complete description of all flips is known [Kollár-Mori92], but it seems very difficult to obtain a list of all divisorial contractions. One can try to study the MMP in two basic ways:

6.1 (Analysis of the MMP). Starting with a projective variety X , let us run the MMP. We obtain a sequence of birational maps

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_i \dashrightarrow X_{i+1} \dashrightarrow \cdots \dashrightarrow X^*.$$

Assume that X has some nice property that we would like to preserve. We need some way of proving that X^* also has this property, at least under some additional assumptions. One way is to prove this directly, by analyzing each step of the MMP. This would sometimes require knowing each step of the MMP, and even in dimension 3 the list is not yet available. Still there are many results that can be established this way, for instance the existence of the MMP itself. In this approach one starts with a variety X and tries to understand every possible way an MMP can *start* with X . This is oftentimes manageable if X has only mild singularities.

Another way is to look at each step of the MMP backwards. In dimension 3 we have a pretty good description of the possible singularities that arise in the course of an MMP. Thus we can start with a variety Y and try to understand every possible way an MMP can *end* with Y . This also seems rather hard. Even the case when Y is smooth

is not at all understood, but in some other cases this approach has been carried through [Kawamata96]. It seems that this method is easier to apply when X is fairly singular.

The *gateway method* attempts to solve the original problem in an intermediate way. In the above chain of maps there is a smallest index i such that X_i is still “nice” but X_{i+1} is not. Hence $X_i \dashrightarrow X_{i+1}$ is a “gateway” through which the process leaves the set of “nice” varieties. Analysing these “gateways” should be easier since the direct approach tends to work for the nice variety X_i and the backwards method tends to work for more complicated singularities of X_{i+1} .

Once such a list of “gateways” is obtained, it is a matter of checking the list to see if some additional properties ensure that this step does not happen.

One of the simplest examples where these ideas yield a nontrivial result is the following.

Example 6.2. Assume that we want to stay within the class of varieties of index 1. In this case there is only one gateway:

Proposition. *Let $f : X \dashrightarrow X'$ be a step of the 3-dimensional MMP where X has index 1 but X' has higher index. Then f is the contraction of a divisor $E \subset X$ to a point. Furthermore, $E \cong \mathbb{P}^2$, X is smooth along E and E has normal bundle $\mathcal{O}_E(-2)$.*

This result is a special case of [Mori88] and [Cutkosky88], though they did not approach this from the point of view of gateways. A proof along the lines suggested by the gateway method is not hard to construct, but this is not any shorter than the direct proofs.

As a consequence we obtain:

Corollary *Let X be a projective 3-fold with index 1 terminal singularities. Assume that X does not contain any surface $S \subset X$ which admits a birational morphism onto $\mathbb{P}^2$. Then each step of the MMP starting with X is a projective 3-fold with index 1 terminal singularities.*

Unfortunately the above condition needs to be checked for every surface S , even for very singular ones. Thus in practice this does not seem to be a useful observation.

6.3. Our aim is to develop a similar theory for real algebraic threefolds. Thus we have to decide which varieties are “nice” and then describe all possible gateways through which the MMP can leave the class of “nice” varieties.

(4.7) naturally suggest a topological choice: X is “nice” if $X(\mathbb{R})$ or $\overline{X(\mathbb{R})}$ is a 3-manifold, maybe with some additional properties. This

was my first attempt, but I was unable to make it work. The main problem seems to be that, as the computations of [Kollár97b] show, there is basically no relationship between the algebraic complexity of a terminal singularity $0 \in X$ and the topological complexity of its real points $X(\mathbb{R})$.

Eventually I settled at a completely algebraic choice: X is nice if it has index 1 along $X(\mathbb{R})$. There are two main reason for adopting this definition:

1. Most complications of 3-dimensional birational geometry come from the appearance of points of index > 1 . Hence this is likely to be the right choice algebraically.
2. One of the first things I realized was that under this condition there would be no flips. Indeed, flips need higher index singular points to exist. If we have only index 1 points along $X(\mathbb{R})$, then all higher index points appear in conjugate pairs. A look at the list of flips [Kollár-Mori92] shows that the singularities appearing along a flipping curve are *always* asymmetrical.

Thus our task is to get a list of all steps $f : Y \rightarrow X$ of the MMP over $\mathbb{R}$ such that Y has index 1 along $Y(\mathbb{R})$. The case of divisor-to-curve contraction is relatively easy. Most of the work is devoted to studying the divisor-to-point contractions. Let $0 \in X(\mathbb{R})$ be the point in question. The existence of f is local in the Euclidean topology. I will go through the classification (up to real analytic equivalence) of 3-dimensional terminal singularities over $\mathbb{R}$ and for each describe all possible $f : Y \rightarrow X$.

There is one subtle point here: the condition of $\mathbb{Q}$ -factoriality is not preserved under analytic equivalence. Thus first we need to develop a notion of “extremal contraction without $\mathbb{Q}$ -factoriality”.

Definition 6.4. Let X be a normal variety over a field K such that K_X is $\mathbb{Q}$ -Cartier. A proper birational morphism $f : Y \rightarrow X$ is called an *elementary extraction* of X if

1. Y is normal and K_Y is $\mathbb{Q}$ -Cartier.
2. The exceptional set $\text{Ex}(f)$ contains a unique K -irreducible divisor E .
3. $-K_Y$ is f -ample.

If we start with Y and construct $f : Y \rightarrow X$ then f is usually called an *elementary contraction* of Y .

We can write $K_Y \equiv f^*K_X + a(E, X)E$ where $a(E, X)$ is the discrepancy of E . Thus $-a(E, X)E$ is f -ample. An exceptional divisor

can never be relatively ample (or nef) (cf. [Kollár-Mori98, 3.35]), thus $a(E, X) > 0$ and so $-E$ is f -ample. This implies that $\text{Ex}(f) = \text{Supp } E$. $f(E)$ is also called the *center* of f on X .

A crucial property of elementary extractions is that they are determined by their exceptional divisors:

Proposition 6.5. *Let X be a normal variety over a field K such that K_X is $\mathbb{Q}$ -Cartier. Let $f_i : Y_i \rightarrow X$ be elementary extractions with exceptional divisors $E_i \subset Y_i$ for $i = 1, 2$. Assume that E_1 and E_2 correspond to each other under the birational map $f_2^{-1} \circ f_1 : Y_1 \dashrightarrow Y_2$. Then Y_1 and Y_2 are isomorphic (over X).*

Proof. Let $\phi : f_2^{-1} \circ f_1 : Y_1 \dashrightarrow Y_2$ be the composition. ϕ is birational, and $\text{Ex}(\phi), \text{Ex}(\phi^{-1})$ have codimension at least 2. Furthermore, K_{Y_1} and $K_{Y_2} = \phi_*(K_{Y_1})$ are relatively ample. Thus ϕ is an isomorphism by an argument of [Matsusaka-Mumford64, p.671]. $\square$

In some sense this gives a way of enumerating all elementary extractions of X . We try to list all exceptional divisors over X and for each construct the corresponding unique elementary extraction. Usually there are infinitely many elementary extractions for a given X and there does not seem to be an easy way to predict for which divisors does the corresponding elementary extraction exist.

The next definition singles out a special class of elementary extractions, by restricting the singularities allowed on Y . The aim is to formalize a special case of the gateway method: we assume that Y is “nice”.

Definition 6.6. Let X be a normal variety over a field K such that K_X is $\mathbb{Q}$ -Cartier. A proper birational morphism $f : Y \rightarrow X$ is called a *gateway-extraction* or *g-extraction* if

1. f is an elementary extraction with exceptional divisor $E \subset Y$.
2. Y has terminal singularities.
3. K_X and E are Cartier at the generic point of every geometrically irreducible K -subvariety of $\text{Ex}(f)$.

In dimension three Y has only isolated singularities, hence (3) is equivalent to the apparently weaker condition:

- 3' K_X and E are Cartier at every K -point of $\text{Ex}(f)$.

If we start with Y and construct $f : Y \rightarrow X$ then f is usually called a *g-contraction* of Y .

The main technical aim of this article is to obtain a list of g -extractions for threefolds with terminal singularities. The project turns out to be

feasible since the discrepancy $a(E, X)$ is always quite small. I have no a priori proof of this, but in every case the study of low discrepancy divisors leads to a description of all g -extractions.

The relationship between low discrepancy divisors and g -extractions rests on the following easy observation:

Proposition 6.7. *Let X be a normal variety over a field K such that K_X is $\mathbb{Q}$ -Cartier. Let $f : Y \rightarrow X$ be a g -extraction with exceptional divisor $F \subset Y$. Let E be a geometrically irreducible K -divisor over X such that $\text{center}_X E \subset \text{center}_X F$. Then*

$$a(E, X) \geq a(E, Y) + a(F, X).$$

Proof. Let $g : Z \rightarrow X$ be a proper birational morphism such that $\text{center}_Z E$ is a divisor on Z . We may assume that the induced rational map $h : Z \dashrightarrow Y$ is a morphism. $h(E)$ is a geometrically irreducible K -subvariety of Y which is contained in $\text{Ex}(f)$. Write

$$\begin{aligned} K_Z &\equiv g^*K_X + a(E, X)E + (\text{other exceptional divisors}), \\ K_Z &\equiv h^*K_Y + a(E, Y)E + (\text{other exceptional divisors}), \\ K_Y &\equiv h^*K_X + a(F, Y)F, \quad \text{and} \\ h^*F &\equiv cE + (\text{other exceptional divisors}), \end{aligned}$$

where $c > 0$ since $h(E) \subset \text{Ex}(f) = \text{Supp } F$ and c is an integer by (6.6.3). Making the substitutions we obtain that $a(E, X) = a(E, Y) + c \cdot a(F, X) \geq a(E, Y) + a(F, X)$. $\square$

The same method also proves the following result:

Proposition 6.8. *Let X be a normal variety over a field K such that K_X is $\mathbb{Q}$ -Cartier. Let $f : Y \rightarrow X$ be a morphism with exceptional divisor $F = \cup F_i \subset Y$. Assume that Y has terminal singularities and K_X and F are Cartier at the generic point of every geometrically irreducible K -subvariety of $\text{Ex}(f)$.*

Let E be a geometrically irreducible K -divisor over X such that $\text{center}_X E \subset \cup_i \text{center}_X F_i$. Then

$$a(E, X) \geq a(E, Y) + \min_i \{a(F_i, X)\}. \quad \square$$

Corollary 6.9. *Let X be a normal variety over a field K such that K_X is $\mathbb{Q}$ -Cartier. Let $f : Y \rightarrow X$ be an elementary extraction with exceptional divisor $E \subset Y$. Assume that E is geometrically irreducible and $a(E, X) \leq 1$.*

Then either $f : Y \rightarrow X$ is a g -extraction, or X has no g -extractions whose center contains $f(E)$.

Proof. Let $g : Z \rightarrow X$ be a g -extraction of X whose center contains $f(E)$. Let $F \subset Z$ be the exceptional divisor. Then $a(E, X) \geq a(E, Z) + a(F, X)$. If $a(E, Z) = 0$ then $\text{center}_Z E$ is a divisor which is contained in F . Since F is an irreducible divisor, $\text{center}_Z E = F$, hence $Y = Z$ by (6.5). Otherwise $a(E, Y) \geq 1$ which would force $a(F, X) \leq 0$. This contradicts (6.6.2). $\square$

Remark 6.10. This corollary gives a very efficient way of finding all g -extractions of a given X in some cases. We have to find *one* geometrically irreducible divisor E such that $a(E, X) \leq 1$ and construct the corresponding elementary extraction $f : Y \rightarrow X$. Then it is usually easy to determine the singularities of Y .

[Markushevich96] proved that if $0 \in X$ is a terminal threefold singularity which is not smooth, then there is a divisor E over $\bar{K}$ with $\text{center}_X E = \{0\}$ and $a(E, X) \leq 1$. Thus there is always such an irreducible K -divisor, but it may not be geometrically irreducible. Still, in many cases we are able to apply (6.9) directly.

In the remaining cases we show that there is always a geometrically irreducible divisor E with $\text{center}_X E = \{0\}$ and $a(E, X) \leq 3$. This is still very useful, thanks to the following:

Corollary 6.11. *Let X be a normal variety over a field K such that K_X is Cartier. Let $f : Y \rightarrow X$ be an elementary extraction with exceptional divisor $E \subset Y$. Assume that E is geometrically irreducible. Let $g : Z \rightarrow X$ be any g -extraction with exceptional divisor F whose center contains $f(E)$.*

Then either $g = f$ or $a(F, X) \leq a(E, X) - 1$.

Proof. By (6.7), $a(E, X) \geq a(E, Z) + a(F, X)$. If $a(E, Z) = 0$ then E and F correspond to each other, hence $Y = Z$ by (6.5). Otherwise $a(E, Y) \geq 1$, thus $a(F, X) \leq a(E, X) - 1$. $\square$

7. SMALL AND DIVISOR-TO-CURVE CONTRACTIONS

In this section we look at those steps $f : X \rightarrow Y$ of the MMP over $\mathbb{R}$ which are either small contractions or contract a divisor to a curve. The two cases can be treated together in the following setting:

Notation 7.1. Let K be a field of characteristic 0. Let X be a 3-fold over K with terminal singularities and $f : X \rightarrow Y$ a proper birational morphism over K such that $-K_X$ is f -ample and $f_*\mathcal{O}_X = \mathcal{O}_Y$. Let $0 \in Y(K)$ be a closed point such that $\dim f^{-1}(0) = 1$.

We will need that under these assumptions $R^1 f_*\mathcal{O}_X = R^1 f_*\mathcal{O}_X(K_X) = 0$ by the generalized Grauert–Riemenschneider vanishing theorem (see, for instance, [CKM88, 8.8] or [Kollár-Mori98, 2.65]).

In keeping with the principles of the gateway method, we are interested in the case when X has index 1 at all points of $X(K)$. The following theorem gives a complete description of such contractions:

Theorem 7.2. *Notation and assumptions as in (7.1). Assume in addition that X has index 1 at all points of $X(K)$. Then Y is smooth at 0 and one can choose local (analytic or formal) coordinates (x, y, z) at $0 \in Y$ such that X is the blow up of the curve $(z = g(x, y) = 0) \subset Y$ for some $g \in K[[x, y]]$.*

In particular, f can not be small.

This theorem has some very useful consequences for the MMP over $\mathbb{R}$:

Corollary 7.3. *Starting with a projective variety X over $\mathbb{R}$, let*

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_i \xrightarrow{f_i} X_{i+1}$$

be the beginning of an MMP over $\mathbb{R}$. Assume that X_j has index 1 at all points of $X_j(\mathbb{R})$ for $j \leq i$. Then the induced maps between the sets of real points

$$X(\mathbb{R}) = X_0(\mathbb{R}) \rightarrow \cdots \rightarrow X_i(\mathbb{R}) \xrightarrow{f_i} X_{i+1}(\mathbb{R})$$

are everywhere defined.

Proof. The only steps of the MMP over $\mathbb{R}$ which are not everywhere defined are the flips of small contractions (4.1). By (7.2) there are no flips in the sequence. $\square$

The topological behavior of divisor-to-curve contractions can also be determined using (7.2):

Theorem 7.4. *Let X be a proper 3-fold over $\mathbb{R}$ with terminal singularities such that X has index 1 at all points of $X(\mathbb{R})$ and $\overline{X(\mathbb{R})}$ is a 3-manifold. Let $f : X \rightarrow Y$ be a proper birational morphism over $\mathbb{R}$ such that $-K_X$ is f -ample and $f_*\mathcal{O}_X = \mathcal{O}_Y$. Assume that $\dim f^{-1}(y) \leq 1$ for every $y \in Y$. Then either*

1. *f is $\mathbb{R}$ -small, or*
2. *$\overline{X(\mathbb{R})}$ contains a 1-sided torus or Klein bottle with nonorientable neighborhood.*

Proof. By (7.2), there is a real curve $D \subset Y$ such that Y is smooth along D and $X = B_D Y$ (at least in a neighborhood of $Y(\mathbb{R})$). Pick $0 \in D(\mathbb{R})$ and let $(z = g(x, y) = 0)$ be a local equation of D . By (7.5), either D is smooth at 0 or X has a unique singular point over 0 with local equation $st = g(x, y)$, which is equivalent to $s^2 - t^2 - g(x, y) = 0$. These are of type $cA_{\geq 1}^-$ or cA_1 in the classification of [Kollár97b]. If g

does not change sign on the (x, y) -plane then $X(\mathbb{R}) \setminus f^{-1}(0) \rightarrow Y \setminus \{0\}$ is one-to-one near 0, hence f is $\mathbb{R}$ -small near 0. If g does change sign on the (x, y) -plane, then from [Kollár97b, sec. 4] we see that (after a coordinate change) $g = \pm(x^2 + y^{2r+1})$ and $X(\mathbb{R})$ is a manifold near $f^{-1}(0)$. In particular, $D(\mathbb{R})$ is the disjoint union of some isolated points and some copies of S^1 .

If $D(\mathbb{R})$ is finite then f is $\mathbb{R}$ -small. Otherwise $D(\mathbb{R})$ has a connected component $M \sim S^1$. Let $E \subset X$ be the exceptional divisor of f . By explicit computation we see that $E(\mathbb{R}) \rightarrow D(\mathbb{R})$ is an S^1 -bundle. Hence there is a unique connected component $N \subset E(\mathbb{R})$ such that N is an S^1 -bundle over M . Thus N is either a torus or a Klein bottle. N is 1-sided with nonorientable neighborhood, since these hold locally for the blow up of a smooth curve in a smooth 3-fold. $\square$

Example 7.5. Set $Y = \mathbb{A}^3$ with coordinates (x, y, z) . Let X be the blow up of the curve $(z = g(x, y) = 0) \subset Y$. Then X has a unique singular point which is given by an equation $st - g(x, y) = 0$.

Corollary 7.6. *Starting with a projective variety X over $\mathbb{R}$, let*

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_i \xrightarrow{f_i} X_{i+1}$$

be the beginning of an MMP over $\mathbb{R}$. Assume that

1. X_j has index 1 at all points of $X_j(\mathbb{R})$ for $j \leq i$,
2. $\overline{X_j(\mathbb{R})}$ is a PL-manifold for $j \leq i$,
3. $\overline{X(\mathbb{R})}$ satisfies the conditions (1.6).

Then:

4. *The induced maps between the sets of real points $f_j : X_j(\mathbb{R}) \rightarrow X_{j+1}(\mathbb{R})$ are everywhere defined for $j \leq i$,*
5. *For every $j \leq i + 1$, there is a finite set $S_j \subset X_j(\mathbb{R})$ such that $\overline{X_j(\mathbb{R})} \setminus S_j$ is homeomorphic to an open subset of $\overline{X(\mathbb{R})}$,*
6. *The smooth part of $\overline{X_{i+1}(\mathbb{R})}$ also satisfies the conditions (1.6).*

Proof. The steps of an MMP are everywhere defined by (7.3).

If $g : U \rightarrow V$ is any divisorial contraction over $\mathbb{R}$ then $\text{Ex}(g^{-1})(\mathbb{R})$ is finite unless g is a divisor-to-curve contraction which is not $\mathbb{R}$ -small.

Let f_j be the first divisor-to-curve contraction in the sequence which is not $\mathbb{R}$ -small. By the above remark, (5) holds for j . By (7.4), $X_j(\mathbb{R})$ contains a surface F which is either a 1-sided torus or Klein bottle with nonorientable neighborhood. We can move F away from any finitely many points, thus by (5) $X(\mathbb{R})$ also contains a 1-sided torus or Klein bottle with nonorientable neighborhood. This is a contradiction. Hence

among the steps there is no divisor-to-curve contraction which is not $\mathbb{R}$ -small. This gives (5) and (6). $\square$

The proof of (7.2) relies on two results:

Proposition 7.7. [Cutkosky88, Thm. 4] *(7.2) holds if K is algebraically closed and C is irreducible.* $\square$

Lemma 7.8. (cf. [Mori88, 1.14]) *Let $f : X \rightarrow Y$ be a proper morphism and $0 \in Y$ a closed point such that $\dim f^{-1}(0) = 1$. Set $\text{red } f^{-1}(0) = C = \cup C_i$.*

1. *If $R^1 f_* \mathcal{O}_X = 0$ then C is a tree of smooth rational curves.*
2. *Let D be a $\mathbb{Q}$ -Cartier Weil divisor on X such that D is Cartier at all but finitely many points of $f^{-1}(0)$. Assume that $(D \cdot C_i) < 0$ for every i and $R^1 f_* \mathcal{O}_X(D) = 0$. Then $-1 \leq (D \cdot C_i) < 0$ for every i and D is not Cartier at the singular points of C .*

Proof. By replacing Y with a neighborhood of 0, we may assume that every fiber of f has dimension at most 1.

Let G be a sheaf on X such that $R^1 f_* G = 0$ and $Q = G/F$ a quotient of G whose support is in $f^{-1}(0)$. We get an exact sequence

$$R^1 f_* G \rightarrow R^1 f_* Q \rightarrow R^2 f_* F.$$

The left hand side is zero by assumption and the right hand side is zero since every fiber of f has dimension at most 1. Thus $R^1 f_* Q = 0$.

Applying this with $G = \mathcal{O}_X$ and $Q = \mathcal{O}_C$ we conclude that $H^1(C, \mathcal{O}_C) = R^1 f_* \mathcal{O}_C = 0$, hence C is a tree of smooth rational curves. This proves (1).

In order to see the second part, we may assume that the residue field of 0 is algebraically closed. Then a point $P \in C$ is singular iff there are at least 2 irreducible components through P .

$\mathcal{O}_X(D) \otimes \mathcal{O}_{C_i}$ is a rank one locally free sheaf except possibly at the points where D is not Cartier. Let L_i denote its quotient by the torsion subsheaf. Then L_i is an invertible sheaf and we have a surjection $\mathcal{O}_X(D) \rightarrow L_i$. Applying $R^1 f_*$ we obtain as above that $H^1(C_i, L_i) = 0$. Thus $\deg L_i \geq -1$.

On the other hand, for every $m > 0$ we have an injection

$$L_i^m \cong (\mathcal{O}_X(D)^{\otimes m} \otimes \mathcal{O}_{C_i}) / (\text{torsion}) \hookrightarrow (\mathcal{O}_X(mD) \otimes \mathcal{O}_{C_i}) / (\text{torsion}).$$

If mD is Cartier then the right hand side has negative degree, thus L_i^m has negative degree. Therefore $\deg L_i = -1$ for every i . Furthermore, $m(D \cdot C_i) \geq m \deg L_i = -m$, so $(D \cdot C_i) \geq -1$.

Set $M := (\mathcal{O}_X(D) \otimes \mathcal{O}_C)/\text{torsion}$. $H^1(C, M) = 0$ as above. We have an exact sequence

$$0 \rightarrow M \rightarrow \sum L_i \rightarrow Q \rightarrow 0,$$

where Q is supported at the singular points of C . Taking cohomologies, we conclude that $H^0(C, Q) = 0$ thus $Q = 0$.

If D is Cartier at a singular point P of C then M is locally free at P and $M \rightarrow \sum L_i$ can not be surjective at P (it is not even surjective when tensored with the residue field at P). $\square$

Corollary 7.9. *Notation and assumptions as in (7.1). Then C is a tree of smooth rational curves and K_X is not Cartier at the singular points of C .*

Proof. Apply (7.8) with $D = K_X$. $R^1 f_* \mathcal{O}_X = R^1 f_* \mathcal{O}_X(K_X) = 0$ by (7.1). $\square$

7.10 (Proof of (7.2)). The assumptions and conclusions are local near 0, thus we may replace Y by a suitable analytic or formal neighborhood of 0.

By (7.9), C is a connected tree of smooth rational curves. $\text{Gal}(\bar{K}/K)$ acts on C , thus C either has a singular K -point or a geometrically irreducible component defined over K .

If $P \in C$ is a singular point then K_X is not Cartier at P by (7.9), but if $P \in X(K)$ then K_X is Cartier at P by assumption. Thus C can not have a singular K -point.

Let $C_0 \subset C$ be a geometrically irreducible component defined over K . Let $H \subset X$ be a divisor defined over K which intersects all irreducible components of $C \setminus C_0$ transversally but is disjoint from C_0 . A large multiple of H defines a morphism $X \rightarrow Y' \rightarrow Y$ such that C_0 is contracted to a point in Y' . If (7.2) holds for $X \rightarrow Y'$, then X has index one along C_0 . By (7.9) this implies that C_0 is a connected component of C . On the other hand, C is connected since $f_* \mathcal{O}_X = \mathcal{O}_Y$. Thus $C = C_0$ and $Y' = Y$.

Therefore it is sufficient to prove (7.2) under the additional assumption that C is geometrically irreducible.

First we show that (7.2) holds if X has only index 1 points along C . By (7.7), Y is smooth at 0 and $X = B_D Y$ where $D \subset Y$ is a curve of embedding dimension 2. D is the image of the exceptional divisor of f , hence D is defined over K . Since D has embedding dimension 2, its ideal is of the form $(z, g(x, y))$.

Finally we show that X has only index 1 points along C . We start with the case when $K = \mathbb{R}$. Let $P_1, \bar{P}_1, \dots, P_k, \bar{P}_k$ be all the conjugate pairs of points of index > 1 . At each P_i pick a local member $D_i \in |K_X|$

such that $C \cap D_i = P_i$. (In order to do this, we may need to replace $X_{\bar{K}}$ with a smaller analytic neighborhood of C .) Let $\bar{D}_i$ be the conjugates. Set $D = \sum D_i$. Let $m > 1$ be the smallest natural number such that mD is Cartier. $D - \bar{D}$ is a Weil divisor and $\mathcal{O}_X(m(D - \bar{D})) \cong \mathcal{O}_X$ since the Picard group of a neighborhood of C is isomorphic to $H^2(C(\mathbb{C}), \mathbb{Z})$ (cf. [Kollár-Mori98, 4.13]).

Corresponding to $1 \in H^0(X, \mathcal{O}_X)$ we obtain an m -sheeted cyclic cover $\pi : \tilde{X} \rightarrow X$ which is unramified outside the points of index > 1 . Thus $K_{\tilde{X}} = \pi^* K_X$ and $\tilde{X}$ has index 1 terminal singularities. Let $\tilde{f} : \tilde{X} \rightarrow \tilde{Y}$ be the Stein factorization of $\tilde{X} \rightarrow Y$. By the already discussed index 1 case, $\tilde{Y}$ is smooth and one can choose local analytic coordinates (x, y, z) at $0 \in \tilde{Y}$ such that $\tilde{X}$ is the blow up of the curve $(z = g(x, y) = 0) \subset \tilde{Y}$.

The group of m^{th} roots of unity (denoted by $\mathbb{Z}_m$) acts on $\tilde{f} : \tilde{X} \rightarrow \tilde{Y}$ and the quotient is $f : X \rightarrow Y$. If $\text{mult}_0 g \geq 2$ then $\tilde{X}$ has a unique singular point (7.5), which is necessarily fixed by the $\mathbb{Z}_m$ -action. Thus X would have a unique point (of index m) which is the quotient of a singular point. On the other hand, the index > 1 singularities of X come in conjugate pairs. Therefore $\tilde{X}$ is smooth and $\tilde{f} : \tilde{X} \rightarrow \tilde{Y}$ is the blow up of a smooth curve $(z = y = 0) \subset \tilde{Y}$.

We can choose local coordinates (x, y, z) on $\tilde{Y}$ such that the action is

$$(x, y, z) \mapsto (\epsilon^a x, \epsilon^b y, \epsilon^c z)$$

where ϵ is a primitive m^{th} root of unity. The corresponding action on $\tilde{X}$ has two fixed points (or a fixed curve) and the corresponding quotients are

$$\mathbb{C}^3 / \frac{1}{m}(a, b - c, c) \quad \text{and} \quad \mathbb{C}^3 / \frac{1}{m}(a, b, c - b).$$

These are both of type cA_0/n on the list (3.15). A simple checking shows that both of these can not be simultaneously terminal.

If K is arbitrary, we can still proceed as above if we can find local divisors $D_i \in |K_X|$ at the index > 1 points such that $(C \cdot \sum D_i) = 0$. Finding the D_i needs a little case by case analysis, and sometimes it can be done only after first taking an auxiliary cover. It is probably easier to observe that there can be at most 2 points of index > 1 along C (see, for instance, [CKM88, 14.5.5]), thus in fact the only case we need to handle is when there is precisely one pair of conjugate points of index > 1 . $\square$

8. PROOF OF THE MAIN THEOREMS

The determination of all divisor-to-point g -extractions is rather technical and lengthy. In this section I state a summary of the list of all g -extractions, and then use it to prove the main theorems stated in the introduction. The proofs of (8.2) and of (8.4) are given in sections 9–11.

Notation 8.1. Let $g(x_1, \dots, x_m)$ be a polynomial or power series and let M be a monomial in the x_i . $M \in g$ means that M appears in g with nonzero coefficient.

Theorem 8.2. *Let $0 \in X$ be a three dimensional terminal singularity over $\mathbb{R}$. If X has a g -extraction then $0 \in X$ is one of the following (up to real analytic equivalence near 0).*

1. (cA_0) Smooth point.
2. $(cA_0/2)$ Quotient of a smooth point by the $\mathbb{Z}_2$ -action $(x, y, z) \mapsto (-x, -y, -z)$.
3. $(cA_{>0}^+)$ Given as $(x^2 + y^2 + g_{\geq m}(z, t) = 0)$, where $g_m(z, t) \neq 0$ and $m \geq 2$.
4. $(cA_{>0}^+/2)$ Given as $(x^2 + y^2 + g_{\geq m}(z, t) = 0)/\mathbb{Z}_2$, where $g_m(z, t) \neq 0$, $m \geq 2$ and the $\mathbb{Z}_2$ -action is $(x, y, z, t) \mapsto (-x, -y, -z, t)$. Furthermore, one of the following two conditions has to be satisfied:
 - (a) m is divisible by 4 and $z^m, t^m \in g$, or
 - (b) m is odd.
5. (cE_6) Given as $(x^2 + y^3 + (z^2 + t^2)^2 + yg_{\geq 4}(z, t) + h_{\geq 6}(z, t) = 0)$.

Complement 8.3. *All the singularities on the above list have g -extractions, with the possible exception of types (4b). These singularities have not been analyzed completely.*

Theorem 8.4. *Let $0 \in X$ be a three dimensional terminal singularity over $\mathbb{R}$ and $f : Y \rightarrow X$ a g -extraction with exceptional divisor $E = \text{red } f^{-1}(0)$. If E is geometrically irreducible then $f : Y \rightarrow X$ is on the following list (up to real analytic equivalence near 0).*

1. $(cA_0, \text{ point blow up})$ $B_0\mathbb{A}^3 \rightarrow \mathbb{A}^3$, $E \cong \mathbb{P}^2$.
2. $(cA_0, \text{ curve blow up})$ $B_C\mathbb{A}^3 \rightarrow \mathbb{A}^3$ where $C \subset \mathbb{A}^3$ is a geometrically irreducible, real and locally planar curve.
3. $(cA_0/2)$ $B_0\mathbb{A}^3/\mathbb{Z}_2 \rightarrow \mathbb{A}^3/\mathbb{Z}_2$, where the $\mathbb{Z}_2$ -action on $\mathbb{A}^3$ is $(x, y, z) \mapsto (-x, -y, -z)$. $E \cong \mathbb{P}^2$.
Furthermore, in this case there are no other g -extractions whose center contains the origin.

4. $(cA_{>0}^+, \text{mult}_0 g \text{ even})$ $X = (x^2 + y^2 + g_{\geq 2m}(z, t) = 0)$ where $g_{2m}(z, t) \neq 0$ and $m \geq 1$. $Y = B_{(m, m, 1, 1)}X$ and $E = (x^2 + y^2 + g_{2m}(z, t) = 0) \subset \mathbb{P}^3(m, m, 1, 1)$.
Furthermore, in this case there are no other g -extractions whose center contains the origin.
5. $(cA_{>0}^+, \text{mult}_0 g \text{ odd})$ $X = (x^2 + y^2 + g_{\geq 2m+1}(z, t) = 0)$ where $g_{2m+1}(z, t) \neq 0$ and $m \geq 1$. This case occurs only if there is a linear change of the (z, t) -coordinates such that $t^{2m+1} \in g$ and $z^i t^j \notin g$ for $i + 2j < 4m + 2$. In this coordinate system, $Y = B_{(2m+1, 2m+1, 1, 2)}X$ and $E = (x^2 + y^2 + g_{2m+1}(z, t) = 0) \subset \mathbb{P}^3(2m+1, 2m+1, 1, 2)$.
6. $(cA_{>0}^+/2, \text{mult}_0 g \text{ even})$ $X = (x^2 + y^2 + g_{\geq 2m}(z, t) = 0)/\mathbb{Z}_2$, where $g_{2m}(z, t) \neq 0$, $m \geq 1$ and the $\mathbb{Z}_2$ -action is $(x, y, z, t) \mapsto (-x, -y, -z, t)$. This case occurs only if m is even and $z^{2m}, t^{2m} \in g$. Then $Y = B_{(m, m, 1, 1)}\tilde{X}/\mathbb{Z}_2$ and $E = \tilde{E}/\mathbb{Z}_2$, where $\tilde{X} = (x^2 + y^2 + g_{\geq 2m}(z, t) = 0)$ and $\tilde{E} = (x^2 + y^2 + g_{2m}(z, t) = 0) \subset \mathbb{P}^3(m, m, 1, 1)$.
Furthermore, in this case there are no other g -extractions whose center contains the origin.
7. $(cA_{>0}^+/2, \text{mult}_0 g \text{ odd})$ $X = (x^2 + y^2 + g_{\geq 2m+1}(z, t) = 0)/\mathbb{Z}_2$ where $g_{2m+1}(z, t) \neq 0$, $m \geq 1$ and the $\mathbb{Z}_2$ -action is $(x, y, z, t) \mapsto (-x, -y, -z, t)$. In this case I do not have a complete list.

Corollary 8.5. *Let $0 \in X$ be a three dimensional terminal singularity over $\mathbb{R}$ and $f : Y \rightarrow X$ a g -extraction with exceptional divisor $E = \text{red } f^{-1}(0)$. Assume that we are not in case (8.4.7). If E is geometrically irreducible then E is normal.*

Proof. Equations for E are given in (8.4). $E \cong \mathbb{P}^2$ in the first two cases. In the remaining cases E is (or is the quotient of) a surface of the form

$$F := (x^2 + y^2 + p(z, t) = 0) \subset \mathbb{P}^3(r, r, 1, s).$$

All the singularities of F are contained in the $(x = y = 0)$ line. Thus we get only finitely many singularities if p is not identically zero, which is always the case in (8.4). $\square$

8.6 (Proof of (8.2) and (8.4) $\Rightarrow$ (1.9) and (1.12)).

Under the additional assumption that $X(\mathbb{R})$ satisfies the conditions (1.6), we need to exclude the cases (8.2.2), (8.2.4), (8.4.2) and in (8.2.3) we need to show that g is not everywhere negative in a punctured neighborhood of 0.

Starting with X , let us run the MMP over $\mathbb{R}$. We get a sequence

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_i \xrightarrow{f_i} X_{i+1}.$$

Assume by induction that (1.9) holds for X_j for $j \leq i$ and (1.12) holds for $f_j : X_j \dashrightarrow X_{j+1}$ for $j \leq i-1$.

We need to show that (1.9) holds for X_{i+1} and (1.12) holds for $f_i : X_i \dashrightarrow X_{i+1}$.

By (7.6) $f_j : X_j(\mathbb{R}) \rightarrow X_{j+1}(\mathbb{R})$ are everywhere defined for $j \leq i-1$ and $\overline{X_i(\mathbb{R})}$ does not contain a 1-sided torus or Klein bottle with nonorientable neighborhood. Furthermore, by (7.4), f_i is either $\mathbb{R}$ -small or a divisor-to-point contraction.

By induction X_i has index 1 along $X_i(\mathbb{R})$, thus f_i is a g-extraction, hence it is one of the cases listed in (8.2). We excluded several cases one at a time.

Excluding (8.2.2). By (8.4.3) $X_i \rightarrow X_{i+1}$ is the blow up of the singular point $\mathbb{A}^3/\mathbb{Z}_2$. This gives a 1-sided $\mathbb{RP}^2$ in $X_i(\mathbb{R})$. This is a contradiction by (7.6).

Excluding (8.4.2). In this case X_i contains a 1-sided torus or Klein bottle with nonorientable neighborhood by (7.4). This is again a contradiction by (7.6).

Excluding (8.2.3) with $g < 0$. That is, we consider the case when $0 \in X_{i+1}$ is of the form $(x^2 + y^2 + g_{\geq m}(z, t) = 0)$ and g is everywhere negative in a punctured neighborhood of 0. (These are called $cA_{>0}^+(0, -)$ in [Kollár97b].) By [Kollár97b, 4.4] the link of $0 \in X_{i+1}(\mathbb{R})$ is a torus. This gives only a 2-sided torus in $X(\mathbb{R})$ which is allowed. I proceed to prove, however, that we still get a 1-sided torus in $X(\mathbb{R})$ coming from the exceptional divisor of $X(\mathbb{R}) \rightarrow X_{i+1}(\mathbb{R})$. This contradicts (1.6.2).

$m = \text{mult}_0 g(z, t)$ is necessarily even, say $m = 2r$. By (8.4.4) the only g-extraction is the $(r, r, 1, 1)$ -blow up. Thus $X_i \rightarrow X_{i+1}$ is this blow up. We distinguish two cases:

General case: $g_{2r}(z, t)$ is negative on $\mathbb{R}^2 \setminus \{0\}$. The exceptional divisor E of the above g-extraction is the weighted hypersurface

$$E = (x^2 + y^2 + g_{2r}(z, t) = 0) \subset \mathbb{P}(r, r, 1, 1).$$

Its canonical divisor is $K_E = \mathcal{O}_E(-2)$, thus E is orientable. The projection $(x : y : z : t) \mapsto (z : t)$ exhibits E as an S^1 -bundle over $\mathbb{RP}^1$, thus $E \sim S^1 \times S^1$. $L(0 \in X_{i+1}(\mathbb{R}))$ is connected, thus $E(\mathbb{R}) \subset X_i(\mathbb{R})$ is a 1-sided torus. By (7.6), $X(\mathbb{R})$ also contains a 1-sided torus, a contradiction.

Special case: $g_{2r}(z, t)$ is not negative on $\mathbb{R}^2 \setminus \{0\}$. $g_{2r}(z, t)$ is the leading term of $g_{\geq m}(z, t)$, which is negative on $\mathbb{R}^2 \setminus \{0\}$. Thus $g_{2r}(z, t)$ is nonpositive on $\mathbb{R}^2 \setminus \{0\}$.

The t -chart on $X_i \cong B_{(r, r, 1, 1)} X_{i+1}$ is $x_1^2 + y_1^2 + t_1^{-2m} g(z_1 t_1, t_1)$. Set $g'(z_1, t_1) := t_1^{-2m} g(z_1 t_1, t_1)$. Then $g'(z_1, t_1)$ is strictly negative outside the z_1 -axis, and is not identically zero on the z_1 -axis. Thus $g'(z_1, t_1)$ is everywhere nonpositive with only finitely many zeros. Thus at each zero of $g'(z_1, t_1)$, X_i has a singular point of type $cA_{>0}^+(0, -)$. This contradicts the inductive assumption.

Thus X_{i+1} does not contain any $cA_{>0}^+(0, -)$ type points.

Excluding (8.2.4). [Kollár97b, 5.9] shows that the link of a singularity of type $cA^+/2$ contains a connected component homeomorphic to $\mathbb{RP}^2$, except when we can write the singularity as $(x^2 + y^2 + g_{\geq m}(z, t) = 0)/\mathbb{Z}_2$, where the $\mathbb{Z}_2$ -action is $(x, y, z, t) \mapsto (-x, -y, -z, t)$ and g is everywhere negative in a punctured neighborhood of 0.

If the link of a singularity of $X_{i+1}(\mathbb{R})$ contains a connected component homeomorphic to $\mathbb{RP}^2$, then $X_i(\mathbb{R})$ contains a 2-sided $\mathbb{RP}^2$. Hence by (7.6), $X(\mathbb{R})$ also contains a 2-sided $\mathbb{RP}^2$ which is excluded by (1.6.1).

Thus we are reduced to the case $(x^2 + y^2 + g_{\geq m}(z, t) = 0)/\mathbb{Z}_2$, where the $\mathbb{Z}_2$ -action is $(x, y, z, t) \mapsto (-x, -y, -z, t)$ and g is everywhere negative in a punctured neighborhood of 0. As in the previous case, $m = \text{mult}_0 g(z, t)$ is necessarily even, say $m = 2r$ and by (8.4.6) the only g -extraction is the $(r, r, 1, 1)$ -blow up. We proceed to prove that the exceptional divisor of $X_i(\mathbb{R}) \rightarrow X_{i+1}(\mathbb{R})$ contains a 1-sided Klein bottle with nonorientable neighborhood, contradicting (1.6.3). We again distinguish two cases:

General case: $g_{2r}(z, t)$ is negative on $\mathbb{R}^2 \setminus \{0\}$. The exceptional divisor E is the $\mathbb{Z}_2$ quotient of the weighted hypersurface

$$\tilde{E} = (x^2 + y^2 + g_{2r}(z, t) = 0) \subset \mathbb{P}(r, r, 1, 1).$$

We already determined that $\tilde{E}(\mathbb{R})$ is a torus and a choice of orientation is given by $(dy \wedge dz)/x$. The $\mathbb{Z}_2$ -action sends this to $(d(-y) \wedge d(-z))/(-x) = -(dy \wedge dz)/x$, hence $\tilde{E}(\mathbb{R})/\mathbb{Z}_2$ is not orientable. We conclude that one of the connected components of $E(\mathbb{R})$ is a Klein bottle. (There may be other connected components.) The Klein bottle is 1-sided since $\tilde{E}(\mathbb{R})$ is 1-sided. The regular neighborhood is nonorientable since its boundary, the link of $0 \in X(\mathbb{R})$, is again a Klein bottle.

Special case: $g_{2r}(z, t)$ is not negative on $\mathbb{R}^2 \setminus \{0\}$.

The same computation as above shows that this leads to a $cA^+/2$ point of the same type that we started with on X_i , which contradicts the inductive assumption.

Thus we conclude that X_{i+1} does not contain any $cA^+/2$ type points. $\square$

8.7 (Proof of (1.1) and (1.7)).

We follow the steps of an MMP over $\mathbb{R}$, using (1.12). $f_i : X_i(\mathbb{R}) \rightarrow X_{i+1}(\mathbb{R})$ is a homeomorphism in cases (1.12.1–2) while (1.12.3) gives a connected sum with $\mathbb{RP}^3$.

In the cases (1.12.4) the exceptional divisor is normal by (8.5), hence we get various cases of (1.1) by (5.7). $\square$

Example 8.8. Consider the singularity $X := (x^2 + y^2 + z^{2m+1} + t^{4m+2} = 0)$. The $(2m+1, 2m+1, 2, 1)$ blow up $X_1 \rightarrow X$ is a g-extraction which is smooth along the $\mathbb{R}$ -points.

The $(m, m, 1, 1)$ -blow up is another g-extraction with one singular point $(x_1^2 + y_1^2 + z_1^{2m+1}t_1 + t_1^{2m+2} = 0)$ on the t -chart. After the $(m+1, m+1, 1, 1)$ -blow up we obtain a variety $X_2 \rightarrow X$ which is smooth along its $\mathbb{R}$ -points.

These two resolutions are indeed quite different. Using the methods of section 5, we see that $X_1(\mathbb{R}) \sim X(\mathbb{R}) \# \mathbb{RP}^3$ and $X_2(\mathbb{R}) \sim X(\mathbb{R}) \# S^1 \times S^2$.

9. cAx AND cD -TYPE POINTS

In this section we begin to classify g-extractions (6.6) of terminal singularities over any field. The classification of 3-fold terminal singularities over nonclosed fields is done in [Kollár97b]. The results are summarized in (3.15). We work through the list of the singularities. In most cases it is easy to see that there are no g-extractions. This is done by exhibiting an elementary extraction (6.4) which is not a g-extraction. If the discrepancy of the exceptional divisor is ≤ 1 then there are no g-extractions by (6.9).

In this section we deal with the cases $cAx/2, cAx/4, cD, cD/2, cD/3$. Among terminal singularities these are somewhat esoteric but the proofs work well for them: in each case (6.9) applies.

The remaining terminal singularities are considered in the next 2 sections. In some cases much more complicated arguments are needed to classify all g-extractions.

Definition 9.1. [Weighted blow-ups]

Let $x_1, \dots, x_n$ be coordinates on $\mathbb{A}^n$. The usual blow up of the origin is patched together from affine charts with morphisms of the form

$$x_j = x'_j x'_i \quad \text{if } j \neq i \quad \text{and} \quad x_i = x'_i.$$

I refer to this as the x_i -chart.

Let $a_1, \dots, a_n$ be a sequence of positive integers. For every $1 \leq i \leq n$ we can define a morphism $\Pi_i : \mathbb{A}^n \rightarrow \mathbb{A}^n$ by

$$x_j = x'_j (x'_i)^{a_j} \quad \text{if } j \neq i \quad \text{and} \quad x_i = (x'_i)^{a_i}.$$

This morphism is birational iff $a_i = 1$ and has degree a_i in general. One can easily notice that Π_i is invariant under the action

$$\mathbb{A}^n(x'_1, \dots, x'_n) / \frac{1}{a_i}(-a_1, \dots, -a_{i-1}, 1, -a_{i+1}, \dots, -a_n)$$

and it descend to a birational morphism π_i

$$\Pi_i : \mathbb{A}^n(x'_1, \dots, x'_n) \rightarrow \mathbb{A}^n(x'_1, \dots, x'_n) / \mathbb{Z}_{a_i} \xrightarrow{\pi_i} \mathbb{A}^n(x_1, \dots, x_n).$$

Furthermore, these charts patch together to give a projective morphism

$$\pi : B_{(a_1, \dots, a_n)} \mathbb{A}^n \rightarrow \mathbb{A}^n.$$

This is called the *weighted blow up* of $\mathbb{A}^n$ with weights $a_1, \dots, a_n$.

Notation 9.2. In the proofs in sections 9–11 I use the following conventions.

First I state the name of the singularity X from (3.15) and possibly some other restrictions. Then I write down the normal form of the equation $X = F(x, y, z, t) / \frac{1}{r}(b_x, b_y, b_z, b_t)$. Any restrictions on F are explained in detail here.

Then I specify the weights (a_x, a_y, a_z, a_t) for a weighted blow up and write down the equation of the birational transform of X on one of the charts on the weighted blow up. Before taking quotients, this has the form $t_1^{-m} F(x_1 t_1^{a_x}, y_1 t_1^{a_y}, z_1 t_1^{a_z}, t_1^{a_t})$ if I use the t -chart. This is denoted by $B\tilde{X}$.

I need to take quotient by 2 actions. First is the $\frac{1}{a_t}(-a_x, -a_y, -a_z, 1)$ -action coming from the weighted blow up. Second, the $\frac{1}{r}(b_x, b_y, b_z, b_t)$ -action needs to be lifted to the (x_1, y_1, z_1, t_1) -space. In some cases this lifts as a $\mathbb{Z}_r$ -action but in other cases the actions combine into a $\mathbb{Z}_{(ra_t)}$ -action. The quotient of $B\tilde{X}$ by these 2 actions is a chart on the weighted blow up of X ; it is denoted by BX .

All these can be done in 4 different charts. I chose the chart where the singularities are most visible or the discrepancy computation is the clearest.

Finally I compute the exceptional divisor of the blow up, the singularities of BX and the discrepancy of the exceptional divisor.

9.3 ($cAx/2$).

Normal form: $ax^2 + by^2 + g_{\geq 4}(z, t)/\frac{1}{2}(1, 0, 1, 1)$, where $ab \neq 0$.

Weights for blow-up: $(1, 1, 1, 1)$

t -chart: $ax_1^2 + by_1^2 + t_1^{-2}g_{\geq 4}(z_1t_1, t_1)/\frac{1}{2}(0, 1, 0, 1)$

Exceptional divisor: $(t_1 = ax_1^2 + by_1^2 = 0)$. Over $\bar{K}$ this is reducible and the two irreducible components are $(t_1 = \sqrt{a}x_1 \pm \sqrt{-b}y_1 = 0)$. The $\mathbb{Z}_2$ -action interchanges these two, so on the quotient we get a geometrically irreducible exceptional divisor.

Singularity: The $\mathbb{Z}_2$ -action has a fixed curve on $B\tilde{X}$: the intersection with the $(x_1 = z_1 = 0)$ -plane. Thus we get a curve of nonterminal singularities on BX .

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = t_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

9.4 ($cAx/4$).

Normal form: $ax^2 + by^2 + g_{\geq 2}(z, t)/\frac{1}{4}(1, 3, 1, 2)$, where $ab \neq 0$ and $g_2(0, 1) = 0$ for weight reasons.

Weights for blow-up: $(1, 1, 1, 1)$

t -chart: $ax_1^2 + by_1^2 + t_1^{-2}g_{\geq 2}(z_1t_1, t_1)/\frac{1}{4}(3, 1, 3, 2)$

Exceptional divisor: $\tilde{E} := (t_1 = ax_1^2 + by_1^2 + g_2(1, 0)z_1^2 = 0)$. $\tilde{E}$ is geometrically irreducible if $g_2(1, 0) \neq 0$. If $g_2(1, 0) = 0$, then $\tilde{E}$ is reducible over $\bar{K}$, and the two irreducible components are $(t_1 = \sqrt{a}x_1 \pm \sqrt{-b}y_1 = 0)$. The $\mathbb{Z}_4$ -action interchanges these two, so on the quotient we get a geometrically irreducible exceptional divisor E .

Singularity: The origin is on $B\tilde{X}$ since $g_2(0, 1) = 0$ and it is a fixed point. We get an index 4 point on BX .

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = t_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

9.5 (cD_4 main series).

Normal form: $x^2 + f_{\geq 3}(y, z, t)$, where we assume that $f_3(y, z, t)$ is irreducible over $\bar{K}$.

Weights for blow-up: $(2, 1, 1, 1)$

x -chart: $x_1 + x_1^{-3}f_{\geq 3}(y_1x_1, z_1x_1, t_1x_1)/\frac{1}{2}(1, 1, 1, 1)$

Exceptional divisor: $\tilde{E} := (x_1 = f_3(y_1, z_1, t_1) = 0)$. $\tilde{E}$ is geometrically irreducible by our assumption.

Singularity: The origin is a fixed point on $B\tilde{X}$, hence we get an index 2 point on BX .

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = x_1 \cdot dy_1 \wedge dz_1 \wedge dt_1$, so $a(E, X) = 1$.

9.6 ($cD_4/2$ main series).

Normal form: $x^2 + f_{\geq 3}(y, z, t)/\frac{1}{2}(1, 1, 0, 1)$, where we assume that $f_3(y, z, t)$ is irreducible over $\bar{K}$. However, for weight reasons $z|f_3(y, z, t)$, so this can not happen.

9.7 ($cD/3$).

Normal form: $x^2 + f_{\geq 3}(y, z, t)/\frac{1}{3}(0, 1, 1, 2)$, where $f_3(0, 0, t) \neq 0$. Since this is not a cE point and for weight reasons, also $f_3(y, z, 0) \neq 0$. We can write $f_3 = t^3 + f_3(y, z, 0)$.

Weights for blow-up: $(2, 1, 1, 1)$

x -chart: $x_1 + x_1^{-3}f_{\geq 3}(y_1x_1, z_1x_1, t_1x_1)/\frac{1}{2}(1, 1, 1, 1)$, and then take the $\mathbb{Z}_3$ -action.

Lifting of the $\mathbb{Z}_3$ -action: It lifts to $\frac{1}{6}(3, 5, 5, 1)$.

Exceptional divisor: $\tilde{E} := (x_1 = f_3(y_1, z_1, t_1) = 0)$. $\tilde{E}$ is geometrically irreducible if $f_3(y, z, 0)$ is not a cube. If $f_3(y, z, 0) = -L(y, z)^3$ over $\bar{K}$, then $\tilde{E}$ has three geometrically irreducible components ($x_1 = t_1 - \eta L(y_1, z_1) = 0$) where $\eta^3 = 1$. The $\mathbb{Z}_6$ -action permutes these, so on BX we get a geometrically irreducible exceptional divisor.

Singularity: The origin is a $\mathbb{Z}_6$ -fixed point which has multiplicity 1 on $B\tilde{X}$. BX has a terminal quotient singularity of index 6.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = x_1 \cdot dy_1 \wedge dz_1 \wedge dt_1$, so $a(E, X) = 1$.

9.8 ($cD_{>4}$ and special cD_4).

Normal form: $x^2 + Q_2(y, z, t)z + g_{\geq 4}(y, z, t)$, where $Q_2(y, 0, t) \neq 0$. In the $cD_{>4}$ we always have this form (with $Q_2(y, z, t) = y^2$). In the cD_4 -case we can achieve this form iff $f_3(y, z, t)$ has a simple linear factor over K .

Weights for blow-up: $(2, 1, 2, 1)$

z -chart: $x_1^2 + Q_2(y_1, z_1, t_1) + z_1^{-4}g_{\geq 4}(y_1z_1, z_1^2, t_1z_1)/\frac{1}{2}(0, 1, 1, 1)$

Exceptional divisor: $\tilde{E} := (z_1 = x_1^2 + Q_2(y_1, 0, t_1) + g_4(y_1, 0, t_1) = 0)$. $\tilde{E}$ is geometrically irreducible iff $Q_2(y_1, 0, t_1)$ is not a square over $\bar{K}$ or $g_4(y_1, 0, t_1) \neq 0$. If $Q_2(y_1, 0, t_1) = -L_1(y_1, t_1)^2$ (over $\bar{K}$) and $g_4(y_1, 0, t_1) = 0$ then $\tilde{E}$ is reducible over $\bar{K}$, and the two irreducible components are $(z_1 = x_1 \pm L_1(y_1, t_1) = 0)$. The $\mathbb{Z}_2$ -action interchanges these two, so $E \subset BX$ is geometrically irreducible.

Singularity: The origin is a fixed point on $B\tilde{X}$, hence we get an index 2 point on BX

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = 2z_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

9.9 ($cD_{>4}/2$ and special $cD_4/2$).

Normal form: $x^2 + Q_2(y, z, t)z + g_{\geq 4}(y, z, t)/\frac{1}{2}(1, 1, 0, 1)$.

Weights for blow-up: $(2, 1, 2, 1)$

z -chart: $x_1^2 + Q_2(y_1, z_1, t_1) + z_1^{-4}g_{\geq 4}(y_1z_1, z_1^2, t_1z_1)/\frac{1}{2}(0, 1, 1, 1)$ and then take the $\mathbb{Z}_2$ -action.

Lifting of the $\mathbb{Z}_2$ -action: We get a pair of commuting $\mathbb{Z}_2$ -action on $B\tilde{X}$, given by $\frac{1}{2}(0, 1, 1, 1)$ and $\frac{1}{2}(1, 0, 1, 0)$.

Singularity: The second action has a fixed curve on $B\tilde{X}$, so BX is singular along a curve.

Exceptional divisor and discrepancy: as in the $cD_{>4}$ -case.

9.10 (cD -cases, conclusion).

We have settled all the $cD_{>4}$, $cD/2$ and $cD/3$ cases, they have no g -extractions.

In the cD_4 cases there are no g -extractions if f_3 is irreducible or if it has a simple linear factor over K . The only remaining case is when f_3 is the product of 3 linear factors which are conjugate over K .

This can not happen when $K = \mathbb{R}$, so over $\mathbb{R}$ points of type cD , $cD/2$ and $cD/3$ do not have g -extractions.

The situation is more complicated over fields which do have cubic extensions, as the following example shows. I have not classified all cases.

Example 9.11. Consider $x^2 + y^3 + az^3 + t^6$, where $a \in K$ is not a cube. The exceptional divisor of the $(3, 2, 2, 1)$ -blow up is irreducible and has discrepancy 1. It has three points of index 2 which are conjugate over K , and no other singularities. Hence this is a g -extraction.

Example 9.12. We obtain an interesting example from the equation $x^2 + (y^2 + z^2)z + t^5$. The $(2, 1, 2, 1)$ blow up has terminal singularities (one with index 2). The exceptional divisor E is singular along a curve.

10. cA -TYPE POINTS

In this section we study g -extractions of cA type terminal singularities. The conventions of (8.1), (9.1) and of (9.2) are used throughout.

10.1 (cA_0). (That is, smooth points.)

Normal form: $\mathbb{A}^3$.

The blow up of the origin is smooth with exceptional divisor $E \cong \mathbb{P}^2$. $a(E, X) = 2$, and by (6.7) $a(F, X) \geq 2$ for every exceptional divisor F with center $X F = \{0\}$. Therefore by (6.11), the blow up of the origin is the only g -extraction.

The exceptional divisor is $E \cong \mathbb{P}^2$ with normal bundle $\mathcal{O}_{\mathbb{P}^2}(-1)$.

10.2 (cA_0/n , $n \geq 2$).

Normal form: $\mathbb{A}^3 / \frac{1}{n}(r, -r, 1)$, where $(r, n) = 1$ and $1 \leq r \leq n - 1$.

Weights for blow-up: $(r, n-r, 1)$.

x -chart: $\mathbb{A}^3(x_1, y_1, z_1) / \frac{1}{r}(1, -n, -1)$.

Exceptional divisor: $\tilde{E} := (x_1 = 0)$. Geometrically irreducible and invariant under the $\mathbb{Z}_r$ -action.

Lifting of the $\mathbb{Z}_n$ -action: The $\mathbb{Z}_n$ -action lifts to $\frac{1}{n}(1, 0, 0)$. Its invariants are $x_2 := x_1^n$ and y_1, z_1 . The $\mathbb{Z}_r$ -action descends to the quotient of the $\mathbb{Z}_n$ -action as $\mathbb{A}^3(x_2, y_1, z_1)/\frac{1}{r}(n, -n, -1)$.

Singularity: We obtain an index r point on the x -chart, and similarly an index $n - r$ point on the y -chart.

Discrepancy: $\pi^* dx \wedge dy \wedge dz = rx_1^n dx_1 \wedge dy_1 \wedge dz_1 = \frac{r}{n} x_1 dx_2 \wedge dy_1 \wedge dz_1$. Since $x_1 = x_2^{1/n}$, we obtain that $a(E, X) = 1/n$.

Conclusion: The above blow up is the only possible g-extraction. If $n \geq 3$ then either $r \geq 2$ or $n - r \geq 2$, and we obtain a singular point of index ≥ 2 on BX .

If $r = 2$ then BX is smooth, the exceptional divisor is $E \cong \mathbb{P}^2$ with normal bundle $\mathcal{O}_{\mathbb{P}^2}(-2)$. $BX \rightarrow X$ is the unique g-extraction.

10.3 (cA_1).

Normal form: $ax^2 + by^2 + cz^2 + dt^m$, where $abcd \neq 0$.

Weights for blow-up: $(1, 1, 1, 1)$

t -chart: $ax_1^2 + by_1^2 + cz_1^2 + dt_1^{m-2}$.

Exceptional divisor: $E := (t_1 = ax_1^2 + by_1^2 + cz_1^2 = 0)$ for $m \geq 3$ and $(t_1 = ax_1^2 + by_1^2 + cz_1^2 + d = 0)$ for $m = 2$. E is geometrically irreducible.

Singularity: BX has exactly one singular point for $m \geq 4$, it lies on the t -chart. BX is smooth for $m = 2, 3$.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = t_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

Conclusion: The only g-extraction is this blow up. The singularities can be resolved by repeatedly blowing up the unique singular point.

10.4 ($cA_1/2$).

Normal form: $ax^2 + by^2 + cz^n + dt^m/\frac{1}{2}(1, 1, 1, 0)$, where $abcd \neq 0$ and $\min\{n, m\} = 2$.

Weights for blow-up: $(1, 1, 1, 1)$

z -chart: $ax_1^2 + by_1^2 + cz_1^{n-2} + dt_1^m z_1^{m-2}$.

Exceptional divisor: $\tilde{E} := (z_1 = ax_1^2 + by_1^2 + c = 0)$ for $m \geq 3$, $(z_1 = ax_1^2 + by_1^2 + dt_1^2 = 0)$ for $n \geq 3$ and $(z_1 = ax_1^2 + by_1^2 + c + dt_1^2 = 0)$ for $n = m = 2$. E is geometrically irreducible.

Singularity: The $\frac{1}{2}(1, 1, 1, 0)$ action lifts to a $\frac{1}{2}(0, 0, 1, 1)$ action. Thus we get a fixed curve, where the blow up intersects the plane $(z_1 = t_1 = 0)$.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = z_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

Conclusion: The only possible g-extraction is this blow up. It has nonterminal singularities, so this does not occur.

10.5 (cA_1/n , $n \geq 3$).

Normal form: $xy + cz^{pm} + dt^2/\frac{1}{n}(r, -r, 1, 0)$, where $(r, n) = 1$ and $cd \neq 0$.

Weights for blow-up: $(1,1,1,1)$

z -chart: $x_1y_1 + cz_1^{pm-2} + dt_1^2/\frac{1}{n}(r-1, 1-r, 1, -1)$

Exceptional divisor: $\tilde{E} := (z_1 = x_1y_1 + dt_1^2 = 0)$, it is geometrically irreducible.

Singularity: The $\mathbb{Z}_n$ -action has an isolated fixed point at the origin on $B\tilde{X}$. Thus BX has an index n point.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = z_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

Conclusion: The only possible g -extraction is this blow up. It has a higher index point, so this does not occur.

10.6 $(cA_{>1}^-)$.

Normal form: $xy + g_{\geq 3}(z, t)$.

Weights for blow-up: $(1,1,1,1)$

t -chart: $x_1y_1 + t_1^{-2}g_{\geq 3}(z_1t_1, t_1)$.

Exceptional divisor: $E := (t_1 = x_1y_1 = 0)$. It has two geometrically irreducible components.

Singularity: Not important

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = t_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

Conclusion: At least 2 geometrically irreducible divisors with discrepancy ≤ 1 .

10.7 $(cA_{>1}/n, n \geq 3 \text{ and } cA_{>1}^-/2)$.

Normal form: $xy + g_{\geq 3}(z, t)/\frac{1}{n}(r, -r, 1, 0)$, where $(r, n) = 1$.

Weights for blow-up: $(1,1,1,1)$

t -chart: $x_1y_1 + t_1^{-2}g_{\geq 3}(z_1t_1, t_1)/\frac{1}{n}(r, -r, 1, 0)$

Exceptional divisor: $\tilde{E} := (t_1 = x_1y_1 = 0)$. It is reducible and both irreducible components are geometrically irreducible and invariant under the $\mathbb{Z}_n$ -action.

Singularity: Not important

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = z_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

Conclusion: At least 2 geometrically irreducible divisors with discrepancy ≤ 1 .

10.8 $(cA_{>1}^+, \text{mult}_0 g \text{ even})$.

Normal form: $ax^2 + by^2 + g_{\geq 2m}(z, t)$, where $m \geq 2$, $-ab$ is not a square and $g_{2m} \neq 0$.

Weights for blow-up: $(m, m, 1, 1)$

t -chart: $ax_1^2 + by_1^2 + t_1^{-2m}g_{\geq 2m}(z_1t_1, t_1)$.

Exceptional divisor: $E := (t_1 = ax_1^2 + by_1^2 + g_{2m}(z_1, 1) = 0)$, it is geometrically irreducible.

Singularity: The t -chart on BX is singular only at points P corresponding to the multiple roots of $g_{2m}(z, 1)$. The singularity at P again

has type $cA_{>1}^+$, but the multiplicity of the corresponding $g^P(z_1, t_1)$ is not necessarily even. The z chart is similar.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = t_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

x -chart: $a + by_1^2 + x_1^{-2m} g_{\geq 2m}(z_1 x_1, t_1 x_1) / \frac{1}{m}(1, 0, -1, -1)$.

Singularity: The fixed points of the $\mathbb{Z}_m$ -action are along the y_1 -axis, this intersects $B\tilde{X}$ in two points $(0, \sqrt{-a/b}, 0, 0)$ which are conjugate over K . Thus BX has 2 index m terminal singularities which are conjugate over K . No other new singular points. The y -chart is similar.

Conclusion: The only g -extraction is the above weighted blow up. The exceptional divisor is geometrically irreducible with a pair of conjugate index m -points. The other singular K -points of BX are again of type $cA_{>1}^+$ or cA_1 .

10.9 ($cA_{>1}^+/2$, $\text{mult}_0 g$ even).

Normal form: $ax^2 + by^2 + g_{\geq 2m}(z, t) / \frac{1}{2}(1, 1, 1, 0)$, where $-ab$ is not a square and $g_{2m} \neq 0$.

Weights for blow up: $(m, m, 1, 1)$.

z -chart: $ax_1^2 + by_1^2 + z_1^{-2m} g_{\geq 2m}(z_1, t_1 z_1) / \frac{1}{2}(1 - m, 1 - m, 1, 1)$.

Exceptional divisor: $\tilde{E} := (z_1 = ax_1^2 + by_1^2 + g_{2m}(1, t_1) = 0)$ is geometrically irreducible.

Singularities: If m is odd then $(z_1 = t_1 = 0)$ intersects $B\tilde{X}$ in a fixed curve of the $\mathbb{Z}_2$ -action, thus we get a singular curve on BX .

If m is even, then on the z -chart the only $\mathbb{Z}_2$ -fixed point is the origin. This is not on $B\tilde{X}$ iff $z^{2m} \in g$.

t -chart: $ax_1^2 + by_1^2 + t_1^{-2m} g_{\geq 2m}(z_1 t_1, t_1) / \frac{1}{2}(1, 1, 1, 0)$.

Singularities: On the t -chart the fixed point set is the t_1 -axis. This intersects the exceptional divisor at the origin. This is not on the blow up iff $t^{2m} \in g$.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = t_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

x -chart: $a + by_1^2 + x_1^{-2m} g_{\geq 2m}(z_1 x_1, t_1 x_1) / \frac{1}{m}(1, 0, -1, -1)$ and we also need to take the quotient by the $\mathbb{Z}_2$ -action.

Lifting the $\mathbb{Z}_2$ -action: $a + by_1^2 + x_1^{-2m} g_{\geq 2m}(z_1 x_1, t_1 x_1) / \frac{1}{2m}(1, 0, m - 1, -1)$

Singularities: On the x -chart the fixed point set is the y_1 -axis. This intersects $B\tilde{X}$ at two points $(0, \pm \sqrt{-a/b}, 0, 0)$. We get a conjugate pair of terminal singularities of index $2m$ on BX .

y -chart: Similar to the x -chart.

Conclusion: $ax^2 + by^2 + g_{\geq 2m}(z, t) / \frac{1}{2}(1, 1, 1, 0)$ where $-ab$ is not a square has a g -extraction iff m is even and $z^{2m}, t^{2m} \in g_{2m}(z, t)$. Under these assumptions, the unique g -extraction is the $(m, m, 1, 1)$ -blow up.

10.10 ($cA_{>1}^+$, $\text{mult}_0 g$ odd).

Normal form: $ax^2 + by^2 + g_{\geq 2m+1}(z, t)$, where $m \geq 1$, $-ab$ is not a square and $g_{2m+1} \neq 0$.

Weights for blow-up: $(s, s, 1, 1)$ for $1 \leq s \leq m$, giving $B_s X \rightarrow X$.

t -chart: $ax_1^2 + by_1^2 + t_1^{-2s} g_{\geq 2m+1}(z_1 t_1, t_1)$.

Exceptional divisor: $E := (t_1 = ax_1^2 + by_1^2 = 0)$, it is irreducible over K but geometrically reducible.

Singularity: The exceptional divisor itself has only smooth or normal crossing points, thus $B_s X$ has only cA type points. The $(x_1 = y_1 = 0)$ line is singular if $s < m$ and generically smooth for $s = m$. $B_m X$ is terminal.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = t_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

Divisors with discrepancy 1: Take the $(1, 1, 1, 1)$ -blow up. BX is singular along a line with an A_{2m-2} transversal section. We can blow up the line $(m-1)$ -times. At each time the exceptional divisor is a pair of transversally intersecting planes, thus we have only cA type singularities. After $(m-1)$ blow ups we obtain $g : Y \rightarrow X$ and Y has only isolated cA points, hence terminal. By (6.8), all the exceptional divisors over $0 \in X$ with discrepancy 1 are birational to divisors on Y . They all come in conjugate pairs and have been enumerated by the above $(s, s, 1, 1)$ blow ups.

Conclusion: There is a unique g -extraction whose exceptional divisor has discrepancy 1. It is the $(m, m, 1, 1)$ -blow up $B_m X \rightarrow X$. Its exceptional divisor is geometrically reducible, so we need to look further.

Divisors with discrepancy 2: Let F be a geometrically irreducible exceptional divisor over $0 \in X$ with discrepancy 2. Then $\text{center}_{B_m X} F$ is real. The center can not be the whole $(x_1 = y_1 = 0)$ line or a smooth point on it since both would give $a(F, X) \geq 3$. Thus it is one of the singular points, corresponding to a linear factor of g_{2m+1} .

By a linear change of the z, t -coordinates we may assume that this linear factor is z . Thus $\text{center}_{B_m X} F$ is the origin of the t chart, where $B_m X$ has equation $ax_1^2 + by_1^2 + t_1^{-2m} g_{\geq 2m+1}(z_1 t_1, t_1)$. This is again a $cA_{>1}^+$ type point, ($\text{mult}_0 g$ can be even or odd) and $a(F, B_m X) = 1$. We have already enumerated all these cases, and we know that F is obtained by an $(r, r, 1, 1)$ -blow up. Putting the two steps together, we see that F is obtained from X by an $(m+r, m+r, 2, 1)$ -blow up. Next we compute these.

Normal form: $ax^2 + by^2 + g_{\geq 2m+1}(z, t)$, where $m \geq 1$, $-ab$ is not a square, $g_{2m+1} \neq 0$ and $\text{mult}_0 g(Z^2, T) \geq 2(m+r)$.

Weights for blow-up: $(m+r, m+r, 2, 1)$, giving $B^r X \rightarrow X$.

z -chart: $ax_1^2 + by_1^2 + z_1^{-2(m+r)}g_{\geq 2m+1}(z_1^2, t_1z_1)/\frac{1}{2}(m+r, m+r, 1, 1)$.

Singularity: If $m+r$ is even, then the action has a fixed curve on $B^r\tilde{X}$, so B^rX is not terminal. If $m+r$ is odd and the origin is in $B^r\tilde{X}$, then we get an index 2 point. $z_1^{-2(m+r)}g_{\geq 2m+1}(z_1^2, t_1z_1)$ does not vanish at the origin iff $z^{m+r} \in g_{\geq 2m+1}(z, t)$. This implies that $r \geq m+1$. But $g_{2m+1}(z_1^2, t_1z_1)$ itself is not divisible by z_1^{4m+3} , hence $r = m+1$.

Conclusion: Assume that there is a linear change of the (z, t) -coordinates such that

$$g_{\geq 2m+1}(z, t) = \sum_{2i+j \geq 2m+2r} \gamma_{ij} z^i t^j, \quad \text{and } \gamma_{ij} \neq 0 \text{ for some } 2i+j = 2m+2r.$$

In this coordinate system, the $(m+r, m+r, 2, 1)$ blow-up gives an elementary extraction whose exceptional divisor is geometrically irreducible and has discrepancy 2. Thus the only possible g-extractions are this weighted blow up and the $(m, m, 1, 1)$ blow up found earlier.

The $(m+r, m+r, 2, 1)$ blow up is a g-extraction only in the $r = m+1$ case:

$$g_{\geq 2m+1}(z, t) = \sum_{2i+j \geq 4m+2} \gamma_{ij} z^i t^j, \quad \text{and } \gamma_{2m+1,0} \neq 0.$$

In some cases (cf. (10.11)), we do not have any geometrically irreducible exceptional divisor over $0 \in X$ with discrepancy 2. Then we have to compute further with discrepancy 3. Fortunately, we can stop there.

Divisors with discrepancy 3:

Normal form: $ax^2 + by^2 + g_{\geq 2m+1}(z, t)$, where $m \geq 1$, $-ab$ is not a square and $g_{2m+1} \neq 0$.

Weights for blow-up: $(2m+1, 2m+1, 2, 2)$, giving $Y \rightarrow X$.

z -chart: $ax_1^2 + by_1^2 + z_1^{-4m-2}g_{\geq 2m+1}(z_1^2, t_1z_1^2)/\frac{1}{2}(1, 1, 1, 0)$.

Exceptional divisor: $E := (t_1 = ax_1^2 + by_1 + g_{2m+1}(1, t_1) = 0)$, it is geometrically irreducible.

Singularity: We get an index 2 point corresponding to the linear factors of $g_{2m+1}(z, t)$. Thus over $\mathbb{R}$ there is always an index 2 point.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = 2z_1^3 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 3$.

Final conclusion: These singularities always have g-extractions. One is the $(m, m, 1, 1)$ -blow up. Its exceptional divisor is geometrically reducible. This is the only g-extraction with discrepancy 1.

In some cases after a suitable coordinate change we can also perform the $(2m+1, 2m+1, 2, 1)$ blow up. This is the only g-extraction whose exceptional divisor is geometrically irreducible and has discrepancy 2.

If $g_{2m+1}(z, t)$ has no linear factors over K , then the $(2m+1, 2m+1, 2, 2)$ blow up is a g-extraction whose exceptional divisor is geometrically irreducible and has discrepancy 3. This is the only one such. This case never happens over $\mathbb{R}$.

There may be other g-extractions whose exceptional divisor is geometrically reducible and has discrepancy 2. I have no such examples.

Example 10.11. Consider the singularity $X := (x^2 + y^2 + z^m + t^n = 0)$ for m, n odd and $m+2 \leq n \leq 2m-1$. The above computations show that there is no geometrically irreducible exceptional divisor over $0 \in X$ with discrepancy ≤ 2 .

10.12 ($cA_{\geq 1}^+/2$, $\text{mult}_0 g$ odd).

Normal form: $ax^2 + by^2 + g_{\geq 2m+1}(z, t)/\frac{1}{2}(1, 1, 0, 1)$, where $-ab$ is not a square and $g_{2m+1} \neq 0$.

Weights for blow up: For weight reasons, only even powers of t appear in g . Thus we can define an integer r by $2m+2r = \text{mult}_0 g(Z^2, T)$. $r \leq m+1$ since $g_{2m+1} \neq 0$. We consider the $(m+r, m+r, 2, 1)$ blow up.

z -chart: $ax_1^2 + by_1^2 + z_1^{-2m-2r} g_{\geq 2m+1}(z_1^2, t_1 z_1)/\frac{1}{2}(m+r, m+r, 1, 1)$ and then we have to take the quotient by the $\frac{1}{2}(1, 1, 0, 1)$ -action. This lifts to a $\frac{1}{2}(1, 1, 0, 1)$ -action on $B\tilde{X}$. We get a pair of commuting $\mathbb{Z}_2$ -actions.

Exceptional divisor: $\tilde{E} := (z_1 = ax_1^2 + by_1^2 + \sum_{2i+j=2m+2r} \gamma_{ij} t_1^j = 0)$ is geometrically irreducible.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = 2z_1^2 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(\tilde{E}, X) = 2$.

Singularities: If $m+r$ is even then the $\mathbb{Z}_2 \times \mathbb{Z}_2$ -action is free in codimension one. One of the elements acts by $(0, 0, 1, 1)$, thus we get a singular curve in BX .

If $m+r$ is odd then one of the elements acts by $(0, 0, 1, 0)$. Coordinates on the quotient are given by $x_1, t_1, z_2 = z_1^2, t_1$ and we get the equation

$$ax_1^2 + by_1^2 + z_2^{-m-r} h_{\geq m+r}(z_2, t_1^2 z_2)/\frac{1}{2}(1, 1, 0, 1)$$

where $h(Z, T^2) = g(Z, T)$. At the origin we get a $\mathbb{Z}_2$ -fixed point unless $z^{m+r} \in g$. Thus $r \geq m+1$. On the other hand $r \leq m+1$, thus $r = m+1$. Computing the t -chart shows that BX has an index 2 point unless $t^{4m+2} \in g$.

Discrepancy: From this we see that $a(E, X) = 1/2$ if $m+r$ is odd and $a(E, X) = 2$ if $m+r$ is even.

Conclusion: If $m+r$ is odd then a g-extraction exists iff

$$g_{\geq 2m+1}(z, t) = \sum_{2i+j \geq 4m+2} \gamma_{ij} z^i t^j, \quad \text{and} \quad \gamma_{2m+1,0} \neq 0 \neq \gamma_{0,4m+2}.$$

If this holds then the $(2m + 1, 2m + 1, 2, 1)$ blow-up is the unique g-extractions. It has a geometrically irreducible exceptional divisor with discrepancy $1/2$.

If $m + r$ is even then there may exist g-extractions with discrepancy $1/2$ or 1 . These can be determined by classifying all $\mathbb{Z}_2$ -invariant divisors of discrepancy 1 and pointwise $\mathbb{Z}_2$ -fixed divisors of discrepancy 2 or 3 over $\tilde{X}$. The first task is easy, and we never get any g-extractions this way. The second task is harder and it seems to require separate consideration of about a dozen cases; I have not done all of them. Fortunately, these singularities can be easily excluded in the main theorems using topological considerations.

11. cE -TYPE POINTS

In this section we study g-extractions of cE type terminal singularities. The conventions of (8.1), (9.1) and of (9.2) are used throughout.

11.1 (cE_6 main series).

Normal form: $x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 4}(z, t)$.

Weights for blow-up: $(2, 2, 1, 1)$

y -chart: $x_1^2 + y_1^2 + h_4(z_1, t_1) + y_1\Phi(y_1, z_1, t_1)/\frac{1}{2}(0, 1, 1, 1)$.

Exceptional divisor: $\tilde{E} := (y_1 = x_1^2 + h_4(z_1, t_1) = 0)$. $\tilde{E}$ is geometrically irreducible iff h_4 is not a square over $\tilde{K}$. If $-h_4$ is a square over K , then $\tilde{E}$ has 2 geometrically irreducible components. In the other cases $\tilde{E}$ is irreducible over K but reducible over $\tilde{K}$. Both of the components are fixed by the $\mathbb{Z}_2$ -action, so the same 3 cases happen for E .

Singularity: The origin is a fixed point of the $\mathbb{Z}_2$ -action which is on $B\tilde{X}$. So we get an index 2 point on BX .

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = 2y_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

Conclusion: If $-h_4$ is a square over K then there are 2 geometrically irreducible divisors with discrepancy 1, so no g-extractions. If h_4 is not a square over $\tilde{K}$ then we get an index 2 point, so again there are no g-extractions.

11.2 ($cE/2$).

Normal form: $x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 4}(z, t)/\frac{1}{2}(1, 0, 1, 1)$. By weight considerations $g_3 = 0$ and $h_5 = 0$. $h_4 \neq 0$ since otherwise we would not have a terminal point. This is a $cE_6/2$ point.

Weights for blow-up: $(2, 2, 1, 1)$

y -chart: $x_1^2 + y_1^2 + h_4(z_1, t_1) + y_1\Phi(y_1, z_1, t_1)/\frac{1}{2}(0, 1, 1, 1)$.

Lifting of the $\mathbb{Z}_2$ -action. The $\mathbb{Z}_2$ -action lifts to $\frac{1}{2}(1, 1, 0, 0)$. Thus on $B\tilde{X}$ we have two commuting $\mathbb{Z}_2$ -actions.

Exceptional divisor: $\tilde{E} := (y_1 = x_1^2 + h_4(z_1, t_1) = 0)$. It is geometrically irreducible iff h_4 is not a square over $\bar{K}$. If $h_4 = -Q_2(z_1, t_1)^2$ then $\tilde{E}$ has 2 geometrically irreducible components ($y_1 = x_1 \pm Q_2(z_1, t_1) = 0$). The $\frac{1}{2}(1, 1, 0, 0)$ action interchanges the 2 components, thus $E \subset BX$ is geometrically irreducible.

Singularity: The $\frac{1}{2}(1, 1, 0, 0)$ action has a fixed curve, thus we get a nonterminal singular curve on BX .

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = 2y_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

11.3 (cE_7 main series).

Normal form: $x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 5}(z, t)$.

Weights for blow-up: $(3, 2, 1, 1)$

x -chart: $x_1 + y_1^3 x_1 + y_1 g_3(z_1, t_1) + h_5(z_1, t_1) + x_1 \Phi(y_1, z_1, t_1) / \frac{1}{3}(1, 1, 2, 2)$.

Exceptional divisor: $\tilde{E} := (x_1 = y_1 g_3(z_1, t_1) + h_5(z_1, t_1) = 0)$. It is geometrically irreducible iff g_3 and h_5 have no common factors.

Singularity: The origin is a fixed point of the $\mathbb{Z}_3$ -action which is on $B\tilde{X}$. So we get an index terminal 3 point on BX . In fact, it is the index 3 terminal point $\mathbb{A}^3 / \frac{1}{3}(1, 1, 2)$.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = 2x_1 \cdot dy_1 \wedge dz_1 \wedge dt_1$, so $a(E, X) = 1$.

Conclusion: If g_3 and h_5 have no common factors then E is irreducible and there are no g -extractions.

11.4 (cE with common linear factors).

Normal form:

$$x^2 + y^3 + yzG_2(z, t) + z^2Q_2(z, t) + zH_4(z, t) + yg_{\geq 4}(z, t) + h_{\geq 6}(z, t).$$

The following cases are of this form:

cE_8 : $x^2 + y^3 + yg_{\geq 4}(z, t) + h_{\geq 5}(z, t)$, if h_5 has a linear factor over K , which we can call z .

cE_7 : $x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 5}(z, t)$, if g_3 and h_5 have a common linear factor over K , which we can call z .

cE_6 : $x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 4}(z, t)$ if there is a linear factor over K , which we can call z , such that $z^2 | h_4$, $z | g_3$ and $z | h_5$.

Weights for blow-up: $(3, 2, 2, 1)$

z -chart:

$$\begin{aligned} x_1^2 + y_1^3 &+ y_1 G_2(0, t_1) + Q_2(0, t_1) + H_4(0, t_1) \\ &+ y_1 g_4(0, t_1) + h_6(0, t_1) + z_1 \Phi(y_1, z_1, t_1) / \frac{1}{2}(1, 0, 1, 1). \end{aligned}$$

Exceptional divisor $\tilde{E}$ is geometrically irreducible:

$$(z_1 = x_1^2 + y_1^3 + y_1 G_2(0, t_1) + Q_2(0, t_1) + H_4(0, t_1) + y_1 g_4(0, t_1) + h_6(0, t_1) = 0).$$

Singularity: The origin is a fixed point of the $\mathbb{Z}_2$ -action which is on $B\tilde{X}$. So we get an index 2 point on BX .

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = 2z_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1$.

11.5 (cE_6 with h_4 a square).

Normal form: $x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 4}(z, t)$.

Weights for blow-up: $(1, 1, 1, 1)$

t -chart:

$$x_1^2 + y_1^3 t_1 + y_1 t_1^2 g_3(z_1, 1) + y_1 t_1^3 g_4(z_1, 1) + t_1^2 h_4(z_1, 1) + t_1^3 h_5(z_1, 1) + t_1^4 \Phi(y_1, z_1, t_1).$$

The z -chart is similar.

Exceptional divisor: $E := (t_1 = x_1 = 0)$, and the scheme theoretic exceptional divisor is $2E$.

Singularity: On the t -chart the singular set is the line $L := (x_1 = y_1 = t_1 = 0)$. We determine the singularities along this line. For a fixed value $z_1 = b \in \bar{K}$ we get a cA -point if $h_4(b, 1) \neq 0$. If $h_4(z, 1)$ has a simple root at b then we get a cD -point. If $h_4(z, 1)$ has a multiple root at b then we still get a cD point if $g_3(b, 1) \neq 0$ and a cE -point if $h_5(b, 1) \neq 0$.

Hence, if $b \in K$ and we do not have a cDV point, then h_4 has a multiple linear factor which also divides g_3 and h_5 . This case was settled in (11.4). Assuming that this is not the case, we obtain that BX has cDV points along L .

The z -chart is similar and easy computations show that the x and y -charts are smooth along E .

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = t_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 2$.

First conclusion: BX is not a g -extraction since it has a singular curve. E is geometrically irreducible and $a(E, X) = 2$, thus if $g : Z \rightarrow X$ is a g -extraction with exceptional divisor F then $a(F, X) = 1$ by (6.11).

Computations: Here we determine all divisors F over $0 \in X$ with $a(F, X) = 1$. If $\text{center}_{BX} F$ is not on L then $a(F, X) \geq 3$, and if $\text{center}_{BX} F$ is a point on L then $a(F, X) \geq 2$. Thus if $a(F, X) = 1$ then $\text{center}_{BX} F = L$ and $a(F, BX) = 0$.

Along L the threefold BX has transversal type A_5 whose singularity is resolved by blowing up the line 3-times. By explicit computation we see that only the first of these produces an exceptional divisor F with $a(F, X) = 1$. This is the same divisor that we encountered in the $(2, 2, 1, 1)$ -blow up and so it was already accounted for.

Final conclusion: There is no g -extraction except possibly when there is a $b \in \bar{K} \setminus K$ such that $(z - bt)^2 | h_4, (z - bt) | g_3, (z - bt) | h_5$. In these cases the same divisibilities hold if we replace b by its conjugates over K . Thus b is quadratic over K , a root of $Q_2(z, 1)$. If F is any divisor

over $0 \in X$ with $a(F, X) = 1$ then its center in BX is $(z_1 - b = 0) \in L$ or its conjugate. Thus F is geometrically reducible.

11.6 (cE_6 last case).

Normal form:

$$x^2 + y^3 + cQ_2(z, t)^2 + yL_1(z, t)Q_2(z, t) + C_3(z, t)Q_2(z, t) + yg_{\geq 4}(z, t) + h_{\geq 6}(z, t),$$

where Q_2 is a quadratic form which is irreducible over K and $-c$ is not a square in K . By a coordinate change as in [AGV85, I.12.6] we can bring this to the simpler form

$$x^2 + y^3 + cQ_2(z, t)^2 + yg_{\geq 4}(z, t) + h_{\geq 6}(z, t),$$

though this is not important.

Normal form and topology over $\mathbb{R}$: We can choose Q_2 to be positive definite and diagonalize it. $-c \in \mathbb{R}$ is not a square, so we can choose $c = 1$. Thus we get the normal form

$$x^2 + y^3 + (z^2 + t^2)^2 + yg_{\geq 4}(z, t) + h_{\geq 6}(z, t).$$

By ([Kollár97b, 4.9]) we obtain that $X(\mathbb{R})$ is homeomorphic to $\mathbb{R}^3$.

g-extractions: As we discussed above, all the g-extractions of X have geometrically reducible exceptional divisors.

Construction of g-extractions: It turns out that in these cases there is a g-extraction. By above remark we do not need to know this for certain to understand the topology over $\mathbb{R}$, thus I only outline the construction.

Basic constructions of toric geometry are used without reference; see [Fulton93] for an introduction.

Over $\bar{K}$ we can bring the equation to the form

$$x^2 + y^3 + z^2t^2 + yg_{\geq 4}(z, t) + h_{\geq 6}(z, t).$$

Let e_x, e_y, e_z, e_t be a basis of $\mathbb{R}^4$. Consider the vectors $w_z = \frac{1}{8}(3, 2, 2, 1)$ and $w_t = \frac{1}{8}(3, 2, 1, 2)$. These vectors give a triangulation of the simplex with vertices e_x, e_y, e_z, e_t where the edges are

$$(e_x, w_z), (e_x, w_t), (e_y, w_z), (e_y, w_t), (e_z, w_z), (e_t, w_t).$$

Let us take the corresponding toric blow up. One can check by a routine but tedious computation that all singularities of BX are terminal and we get two index 3 points on the chart corresponding to the simplex (e_x, e_y, w_z, w_t) .

Note that the above construction is symmetric in z and t . Thus if we start with a quadratic form $Q_2 = z^2 + qt^2$ and introduce new coordinates $z' = z + \sqrt{q}t$ and $t' = z - \sqrt{q}t$ then $Q_2 = z't'$ and any blow up which is symmetric in z', t' can be transformed back to a blow up

of X defined over K . We need to check that the two index 3 points become conjugates over K , but this is easy to see from the explicit equations.

11.7 (cE_7 with common nonlinear factor).

Normal form: $x^2 + y^3 + yg_{\geq 3}(z, t) + h_{\geq 5}(z, t)$, where we assume that the greatest common divisor of g_3 and h_5 is K -irreducible (and non-constant). We write $g_3 = Q(z, t)G(z, t)$ and $h_5 = Q(z, t)H(z, t)$. (Q is allowed to be linear, though this case is treated already.)

Weights for blow-up: $(3, 2, 1, 1)$

x -chart: $x_1 + y_1^3 x_1 + y_1 g_3(z_1, t_1) + h_5(z_1, t_1) + x_1 \Phi(y_1, z_1, t_1) / \frac{1}{3}(1, 1, 2, 2)$.

Exceptional divisor: It has two irreducible components:

$$\begin{aligned}\tilde{E} &:= (x_1 = y_1 G(z_1, t_1) + H(z_1, t_1) = 0), \quad \text{and} \\ \tilde{F} &:= (x_1 = Q(z_1, t_1) = 0).\end{aligned}$$

$\tilde{E}$ is geometrically irreducible, $\tilde{F}$ is irreducible but geometrically reducible if Q is not linear.

Discrepancy: $\pi^* \frac{dy \wedge dz \wedge dt}{x} = 3x_1 \frac{dy_1 \wedge dz_1 \wedge dt_1}{x_1}$, so $a(E, X) = 1 = a(F, X)$. (The latter equality uses that Q is not a multiple factor.)

Further aim: We would like to construct a birational morphism $g : Z \rightarrow X$ whose exceptional divisor corresponds to E , and determine the singularities of Z . Thus in BX we have to contract F . F is not $\mathbb{Q}$ -Cartier in BX and F can not be contracted in BX . First we have to correct this problem.

Singularities of BX : I claim that BX has only canonical singularities. This can be done 2 ways. One can compute each chart explicitly, which is rather tedious. I found it easier to use a degeneration argument as follows. Let F be the normal form of the equation as above. We may assume that $g_3(1, 0) = 1$. Consider the substitution

$$F(x, y, z, t) \mapsto \epsilon^{-24} F(\epsilon^{12} x, \epsilon^8 y, \epsilon^6 z, \epsilon^7 t).$$

The exponents are chosen so that for $\epsilon \rightarrow 0$ the limit is $X_0 := (x^2 + y^3 + yz^3 = 0)$. The $(3, 2, 1, 1)$ -blow up BX_0 is easy to compute. We find an index 3 terminal point $\mathbb{A}^3 / \frac{1}{3}(1, 1, 2)$, a curve of cA -points and a curve of cE_7 -points corresponding to the t -axis. Thus BX , as a small deformation of BX_0 , has an index 3 point at the origin and some cDV singularities. (These turn out to be isolated points but we do not need this.) As in (11.3) we see that the index 3 point is at the origin of the x -chart and it is $\mathbb{A}^3 / \frac{1}{3}(1, 1, 2)$. In particular it is $\mathbb{Q}$ -factorial.

Small blow up: Let $p : Y \rightarrow BX$ be the blow up of F in BX . Let $F' \subset Y$ denote the birational transform of F . Away from the index 3 point BX is locally isomorphic to $B\tilde{X}$. $\tilde{F}$ is defined by 2 equations

($x_1 = Q(z_1, t_1) = 0$), thus $p : Y \rightarrow BX$ is small and is an isomorphism at all points where F is $\mathbb{Q}$ -Cartier. The index 3 point is $\mathbb{Q}$ -factorial, so F is $\mathbb{Q}$ -Cartier there. Thus $p : F' \rightarrow F$ is an isomorphism.

Contracting F' : F is a cone over a K -irreducible curve, hence its cone of curves over K is 1-dimensional. If $C \subset F'$ is a general curve then $(C \cdot K_Y) = (p(C) \cdot K_{BX}) < 0$ and $(C \cdot F') = (p(C) \cdot F) < 0$. Thus the curves in F' generate a K_Y -negative extremal ray of Y/X , which can be contracted. We obtain $f : Y \rightarrow Z$ and $g : Z \rightarrow X$. $P := f(F')$ is a K -point since F' is connected.

Conclusion: $g : Z \rightarrow X$ has a geometrically irreducible exceptional divisor corresponding to E and it has discrepancy 1. Furthermore, by (6.11) the index of P can not be one since $a(F, X) = 1$. Hence there are no g -extractions.

11.8 (Conclusion). The cE_8 case is settled if $h_5(z, t)$ has a linear factor over K . This always holds if $K = \mathbb{R}$, hence at least in this case there are no g -extractions. I do not know what happens if $K \neq \mathbb{R}$.

The cE_7 case is settled if $g_3(z, t)$ and $h_5(z, t)$ have no common factor, or if they have a common linear factor over K or if they have a unique common factor over K . This accounts for all the possibilities, hence there are no g -extractions.

The cE_6 case is settled if $h_4(z, t)$ is not a square over $\bar{K}$, if $-h_4(z, t)$ is a square over K or if $h_4(z, t)$ is divisible by the square of a linear form over K . In these cases there are no g -extractions.

The remaining case is treated in (11.6) and the unique g -extraction is written down explicitly. For the applications in this paper the existence is not crucial.

Example 11.9. Let X be the cE_7 type singularity $x^2 + y^3 + yg_3(z, t) + h_5(z, t)$, where g_3 and h_5 do not have a common factor. It is not hard to see that X is an isolated singular point and its $(3, 2, 1, 1)$ -blow up has only terminal singularities. As in (11.3), the y chart on the blow up gives the exceptional divisor

$$E = (g_3(z, t) + h_5(z, t) = 0)/\frac{1}{2}(1, 1, 1, 1).$$

This gives examples of extremal contractions whose exceptional divisor E has a quite complicated singularity along the $(z = t = 0)$ -line.

1. $x^2 + y^3 + yz^3 + t^5$. E is singular along $(z = t = 0)$, with a transversal singularity type $z^3 + t^5$, that is E_8 .
2. $x^2 + y^3 + y(z - at)(z - bt)(z - ct) + t^5$. E has triple selfintersection along $z = t = 0$.

12. HYPERBOLIC 3-MANIFOLDS

The aim of this section is to show that every hyperbolic 3-manifold satisfies the conditions (1.6).

Theorem 12.1. *Let M be a compact hyperbolic 3-manifold. Then M does not contain any PL submanifold of the following types:*

1. $\mathbb{RP}^2$
2. 1-sided $S^1 \times S^1$
3. 1-sided Klein bottle.

We use two facts about hyperbolic 3-manifolds. First, that their universal cover is homeomorphic to $\mathbb{R}^3$. Second, that their fundamental group does not contain a subgroup isomorphic to $\mathbb{Z}^2$ (see, for instance, [Scott83, 4.6]).

More generally, we see how these conditions fit in the framework of Thurston's geometrization conjecture. This version was pointed out to me by Kapovich.

Theorem 12.2. *Let M be a compact 3-manifold. Assume that $M = M_1 \# \cdots \# M_k$, where*

- (i) *each M_i is aspherical, and*
- (ii) *the Seifert fibered part of the Jaco–Shalen–Johannson decomposition of M_i is orientable.*

Then M does not contain any PL submanifold of the following types:

1. $\mathbb{RP}^2$
2. 1-sided $S^1 \times S^1$
3. 1-sided Klein bottle.

We consider the 3 types of submanifolds separately. Condition (1.6.1) is closely related to the notion of $\mathbb{P}^2$ -irreducibility (cf. [Hempel76, p.88]).

Lemma 12.3. *Let M be a 3-manifold with universal cover $\tilde{M}$.*

1. *If $M \sim M_1 \# M_2$, then M contains a 2-sided $\mathbb{RP}^2$ iff one of the summands does.*
2. *Assume that $\tilde{M}$ is homeomorphic to $\mathbb{R}^3$. Then M does not contain an $\mathbb{RP}^2$ and M can not be written as a nontrivial connected sum.*

Proof. Assume that $F \subset M$ is a 2-sided $\mathbb{RP}^2$. We may assume that F is transversal to the gluing S^2 . Thus $C = F \cap S^2$ is an embedded curve in F . Assume first that F has a connected component $C_1 \subset C$ which is not null homotopic in F . Then F is not orientable along C_1 , and the same holds for M along C_1 . But M is orientable along S^2 , a contradiction.

Take any connected component $C_i \subset C$ such that $C_i \subset S^2$ bounds a disc D_i which is disjoint from C . C_i also bounds a disc D'_i in F (since it is null homotopic in F). Thus we can change the embedding $\mathbb{RP}^2 \rightarrow M$ by replacing D'_i with D_i and then pushing it to one side. The new embedding is still 2-sided. Repeating if necessary, we eventually get an embedding which is disjoint from S^2 , proving (1).

$\mathbb{RP}^2$ can not be embedded into $\mathbb{R}^3$ (cf. [Greenberg-Harper81, 27.11]), thus the preimage of $\mathbb{RP}^2$ in $\mathbb{R}^3$ is a union of copies of S^2 . Fix one of these and call it N . By the Schoenflies theorem (cf. [Moise77, Sec. 17]) N bounds a 3-ball B^3 . At least one element of $\pi_1(M)$ maps N to itself. It can not map the inside of N to its outside since these are not homeomorphic. If it maps B^3 to itself, then by the Borsuk–Ulam theorem (cf. [Fulton95, 23.20]) we have a covering transformation with a fixed point, a contradiction.

Assume that we have $S^2 \sim N' \subset M$ and let $S^2 \sim N \subset \tilde{M}$ be one of the preimages. Then N bounds a 3-ball and so does N' . $\square$

In order to study the conditions (1.6.2–3) we have to distinguish two cases.

12.4 (Incompressible case). Let M be a compact 3-manifold and $S \subset M$ a compact 1-sided torus or Klein bottle. Assume that $\pi_1(S) \hookrightarrow \pi_1(M)$. Let ∂U be the boundary of a regular neighborhood of S . Then ∂U is a 2-sided torus or Klein bottle and $\pi_1(\partial U) \hookrightarrow \pi_1(S) \hookrightarrow \pi_1(M)$ is an injection. This implies that ∂U is incompressible in M (cf. [Hempel76, pp.88-89]). Thus U is one of the pieces of the Jaco–Shalen–Johannson decomposition of M (cf. [Scott83, p.483]). We have to be a little more careful since U is Seifert fibered, thus it may sit inside one of the Seifert fibered components.

The fundamental group of a hyperbolic 3-manifold does not contain a subgroup isomorphic to $\mathbb{Z}^2$ (see, for instance, [Scott83, 4.6]), hence the incompressible case does not happen for hyperbolic 3-manifolds.

12.5 (Compressible case). In this case we show that M can be written as a connected sum with a very special summand.

Proposition 12.6. *Let M be a compact 3-manifold. Then M contains a 1-sided torus T such that $\pi_1(T) \rightarrow \pi_1(M)$ is not an injection iff $M \sim N \# (S^1 \tilde{\times} S^2)$ or $M \sim N \# (S^1 \times \mathbb{RP}^2)$*

Proof. Let $T \subset U \subset M$ be a regular neighborhood. Set $V = M \setminus U$. Then $\partial U = \partial V \sim S^1 \times S^1$. We know that $\pi_1(\partial U)$ injects into $\pi_1(U)$. If $\pi_1(\partial U) \hookrightarrow \pi_1(V)$, then $\pi_1(\partial U) \hookrightarrow \pi_1(U) \hookrightarrow \pi_1(M)$ by Schreier’s theorem (cf. [Lyndon-Schupp77, IV.2.6]). $\pi_1(\partial U)$ is an index

2 subgroup of $\pi_1(T)$ and $\pi_1(T)$ is torsion free. Thus $\pi_1(T) \rightarrow \pi_1(M)$ is also an injection, a contradiction.

Therefore, by the Loop theorem (cf. [Hempel76, 4.2]), there is an embedding of the disc $j : (B, \partial B) \hookrightarrow (V, \partial V)$ such that the image of $j(\partial B)$ is not contractible in ∂V .

Let us cut V along $j(B)$ to get W . The boundary of W is ∂V cut along $j(\partial B)$ (which is a cylinder) with two copies of B pasted to the ends. That is, $\partial W \sim S^2$. Therefore M is obtained by pasting W to a 3-manifold (with boundary) K , which is obtained from U by attaching a 2-handle.

There are two cases corresponding to whether $j(\partial B)$ is a primitive element of $\pi_1(U) \cong \mathbb{Z}^2$ (hence $\pi_1(K) \cong \mathbb{Z}$) or is contained in $2\mathbb{Z}^2$ (hence $\pi_1(K) \cong \mathbb{Z} + \mathbb{Z}_2$). $\square$

Proposition 12.7. *Let M be a compact 3-manifold which does not contain a 2-sided $\mathbb{RP}^2$. Then M contains a 1-sided Klein bottle K such that $\pi_1(K) \rightarrow \pi_1(M)$ is not an injection iff $M \sim N \# (S^1 \tilde{\times} S^2)$ or $M \sim N \# (\mathbb{RP}^3 \# \mathbb{RP}^3)$.*

Proof. Let $K \subset U \subset M$ be a regular neighborhood. As in the proof of (12.6) we obtain an embedding of the disc $j : (B, \partial B) \hookrightarrow (V, \partial V)$ such that the image of $j(\partial B)$ is not contractible in ∂V . We again cut V along $j(B)$ to get W . Let ∂V^* denote ∂V cut along $j(\partial B)$.

There are 3 cases to consider corresponding to what ∂V^* is:

1. (∂V^* is a cylinder). Then we obtain a connected sum decomposition as in (12.6).
2. (∂V^* consists of two Moebius bands). Then ∂W is two disjoint projective planes, hence M contains a 2-sided projective plane. This can not happen by assumption.
3. (∂V^* is a Moebius band). In this case $j(\partial B)$ is 1-sided in ∂V , thus M is not orientable along $j(\partial B)$. Then $j(\partial B)$ can not be the boundary of an embedded disc. $\square$

Remark 12.8. So far we have excluded Seifert fiber spaces from considerations. Many Seifert fiber spaces do contain 1-sided tori or Klein bottles.

If $p : M \rightarrow F$ is a Seifert fiber space and $C \subset F$ a 1-sided curve not passing through any critical value, then $p^{-1}(C) \subset M$ is a 1-sided torus or Klein bottle. Another example can be obtained as follows. Let $x, x' \in F$ be two points such that the fibers over them have multiplicity 2. Let $I \subset F$ be a simple path connecting x and x' . Then $p^{-1}(I)$ is a 1-sided Klein bottle.

It is not hard to see that if $T \subset M$ is a 1-sided torus or Klein bottle such that $p(T)$ is 1-dimensional (these are called vertical) then T is obtained by one of the above constructions.

Assume now in addition that M has a geometry modelled on $\mathbb{H}^2 \times \mathbb{R}$ (cf. [Scott83, p.459]). Then by [Johannson79, 5.6], every 1-sided torus or Klein bottle in M is isotopic to a vertical one.

This way we obtain many examples of nonorientable Seifert fiber spaces which satisfy the conditions (1.6).

REFERENCES

- Akbulut-King81. S. Akbulut and H. King, All knots are algebraic, *Comm. Math. Helv.* 56 (1981) 339-351
- Akbulut-King92. S. Akbulut and H. King, *Topology of Real Algebraic Sets*, MSRI Publ. vol.25, Springer 1992
- AGV85. V. I. Arnold, S. M. Gusein-Zade and A. N. Varchenko, *Singularities of Differentiable Maps I-II*, Birkhäuser 1985
- Andreatta-Wiśniewski96. M. Andreatta and J. Wiśniewski, A survey on contractions of higher dimensional varieties, in *Algebraic Geometry*, Santa Cruz 1995, Amer. Math. Soc. 1997
- BCR87. J. Bochnak, M. Coste and M-F. Roy, *Géométrie algébrique réelle*, Springer 1987
- CKM88. H. Clemens, J. Kollár and S. Mori, *Higher Dimensional Complex Geometry*, Astérisque vol.166, 1988
- Kollár-Mori98. J. Kollár and S. Mori, *Birational Geometry of Algebraic Varieties*, Cambridge Univ. Press (to appear) 1998
- Comessatti14. A. Comessatti, Sulla connessione delle superficie razionali reali, *Annali di Math.* 23(3) (1914) 215-283
- Cutkosky88. S. D. Cutkosky, Elementary contractions of Gorenstein threefolds, *Math. Ann.* 280 (1988) 521-525
- Fulton93. W. Fulton, *Introduction to toric varieties*, Princeton Univ. Press, 1993
- Fulton95. W. Fulton, *Algebraic topology*, Springer, 1995
- Greenberg-Harper81. M. Greenberg and J. Harper, *Algebraic topology*, Benjamin/Cummings, 1981
- Hayakawa97. T. Hayakawa, (personal communication)
- Hempel76. J. Hempel, *3-manifolds*, Princeton Univ. Press, 1976
- Iskovskikh80. V. A. Iskovskikh, Anticanonical models of three-dimensional algebraic varieties, *J. Soviet Math.* 13 (1980) 745-814
- Johannson79. K. Johannson, Homotopy equivalences of 3-manifolds with boundaries, *Springer Lecture Notes* vol. 761, 1979
- Kawamata92. Y. Kawamata, Boundedness of $\mathbb{Q}$ -Fano threefolds, *Proc. Int. Conf. Algebra*, *Contemp. Math.* vol. 131 (1992) 439-445
- Kawamata96. Y. Kawamata, Divisorial contractions to 3-dimensional terminal quotient singularities, in *Higher dimensional complex varieties* (Trento, 1994), de Gruyter, (1996) 241-246
- Kharlamov76. V. Kharlamov, The topological type of non-singular surfaces in RP^3 of degree four, *Funct. Anal. Appl.* 10 (1976) 295-305

- Kollár87. J. Kollár, The structure of algebraic threefolds - an introduction to Mori's program, Bull. AMS 17 (1987) 211-273
- Kollár90. J. Kollár, Minimal Models of Algebraic Threefolds: Mori's Program, Astérisque 177-178(1990) 303-326
- Kollár97a. J. Kollár, Real Algebraic Surfaces, Notes of the 1997 Trento summer school lectures, (preprint)
- Kollár97b. J. Kollár, Real Algebraic Threefolds I. Terminal Singularities (preprint)
- Kollár et al.92. J. Kollár (with 14 coauthors), Flips and Abundance for Algebraic Threefolds, Astérisque 211 (1992)
- KoMiMo92. J. Kollár - Y. Miyaoka - S. Mori, Rationally Connected Varieties, J. Alg. Geom. 1 (1992) 429-448
- Kollár-Mori92. J. Kollár - S. Mori, Classification of three dimensional flips, Jour. AMS, 5 (1992) 533-703
- Kollár-Mori98. J. Kollár - S. Mori, Birational geometry of algebraic varieties, Cambridge Univ. Press, 1998 (to appear)
- Lyndon-Schupp77. R. Lyndon and P. Schupp, Combinatorial group theory, Springer, 1977
- Markushevich96. D. G. Markushevich, Minimal discrepancy for a terminal cDV singularity is 1, J. Math. Sci. Univ. Tokyo 3(1996) 445-456
- Matsusaka-Mumford64. T. Matsusaka - D. Mumford, Two fundamental theorems on deformations of polarised varieties, Amer. J. Math. 86 (1964) 668-684
- Moise77. E. Moise, Geometric topology in dimensions 2 and 3, Springer, 1977
- Mori82. S. Mori, Threefolds whose Canonical Bundles are not Numerically Effective, Ann. of Math. 116 (1982) 133-176
- Mori85. S. Mori, On 3-dimensional terminal singularities, Nagoya Math. J. 98 (1985) 43-66
- Mori88. S. Mori, Flip theorem and the existence of minimal models for 3-folds, Journal AMS 1 (1988) 117-253
- Nash52. J. Nash, Real algebraic manifolds, Ann. Math. 56 (1952) 405-421
- Reid80. M. Reid, Canonical Threefolds, in Géométrie Algébrique Angers, A. Beauville ed., Sijthoff & Noordhoff, 1980, pp. 273-310
- Reid85. M. Reid, Young person's guide to canonical singularities, in Algebraic Geometry, Proc. Symp. Pure Math. vol.46, pp. 345-414
- Riesler93. J.-L. Riesler, Construction d'hypersurfaces réelle (Sém. Bourbaki #763), Astérisque, 216 (1993) 69-86
- Rolfsen76. D. Rolfsen, Knots and links, Publish or Perish, 1976
- Scott83. P. Scott, The geometries of 3-manifolds, Bull. London M.S., 15 (1983) 401-487
- Segre42. B. Segre, The non-singular cubic surfaces, Clarendon Press, 1942
- Silhol84. R. Silhol, Real algebraic surfaces with rational or elliptic fibering, Math. Zeitschr. 186(1984) 465-499
- Silhol89. R. Silhol, Real algebraic surfaces, Springer Lecture Notes vol. 1392, 1989
- Viro90. O. Ya. Viro, Real algebraic plane curves, Leningrad Math. J. 1 (1990) 1059-1134

University of Utah, Salt Lake City UT 84112

